


RECLAMATION

Managing Water in the West

Vegetation Response to Interim Flows in the San Joaquin River

Annual Report 2013



U.S. Department of the Interior
Bureau of Reclamation
Technical Service Center
Denver, Colorado

April 2014

Mission Statements

The U.S. Department of the Interior protects America's natural resources and heritage, honors our cultures and tribal communities, and supplies the energy to power our future.

The mission of the Bureau of Reclamation is to manage, develop, and protect water and related resources in an environmentally and economically sound manner in the interest of the American public.

Vegetation Response to Interim Flows in the San Joaquin River

Annual Report 2013

prepared by

Rebecca Siegle, Natural Resources Specialist

Gregory Reed, Natural Resources Specialist

S. Mark Nelson, Research Aquatic Biologist

Technical Service Center, Denver, CO

Erin Rice, Natural Resources Specialist

San Joaquin River Restoration Program

prepared for

**Mid- Pacific Region
Sacramento, California**



**U.S. Department of the Interior
Bureau of Reclamation
Technical Service Center
Denver, Colorado**

April 2014

Contents

	Page
Introduction.....	1
Background	1
Project Area	2
Methods.....	5
Vegetation Transects.....	5
Timing.....	5
Understory Vegetation	5
Overstory Vegetation	8
Stem Density	8
Habitat Variables	8
Statistical Analysis.....	9
Photo Stations	9
Groundwater Monitoring	10
Results	10
Vegetation Transects.....	10
Timing.....	10
Understory Vegetation	10
Overstory Vegetation	14
Stem Density	14
Habitat Variables	17
Statistical Analysis.....	17
Photo Stations	20
Groundwater Monitoring	20
Discussion.....	22
Reach 1A.....	22
Reach 1B.....	22
Reach 2A.....	23
Reach 2B.....	23
Reach 3.....	24
Reach 4A.....	25
East Side Bypass	25
Mariposa Bypass	26
Reach 4B2	27
Reach 5.....	27
Conclusions.....	27
Summary.....	29
Literature Cited	30

Contents, continued

Appendices

Appendix A—Maps of Vegetation Transects by River Reach

Appendix B—Vegetation Transects Waypoints

Appendix C—Data Collection Forms

Appendix D—Scientific Names and Locations of Plants Detected in Vegetation Transects 2011 to 2013

Appendix E— Total percent cover of individual plant species detected in the understory layer of vegetation transects from 2011 to 2013

Appendix F—Photo Stations August 2011 to June 2013

Tables

Page

1	Average total percent cover by type in the understory layer of vegetation transects along the San Joaquin River from 2011 to 2013	11
2	Species richness (number of species detected) for herbaceous and woody plants by reach along the San Joaquin River from 2011 to 2013.....	12
3	Proportion of native and introduced species in the herbaceous and overstory layers by reach along the San Joaquin River from 2011 to 2013	14
4	Total percent cover and average height of woody overstory species (>1 m) detected in vegetation transects for upstream reaches (A) and downstream reaches (B) of the San Joaquin River from 2011 to 2013	15
5	Density of woody plant species by size class and species in river reaches along the San Joaquin River from 2011 to 2013	16
6	Ranking of riparian habitat variables as an indicator of riparian condition by reach along the San Joaquin River from 2011 to 2013	18
7	Statistical results comparing total plant cover and density over time for various parameters in upstream and downstream reaches of the San Joaquin River	20

Figures

Page

1	Location of upstream vegetation transects in Reaches 1, 2, and 3 along the San Joaquin River	6
2	Location of downstream vegetation transects in Reaches 4 and 5, Eastside Bypass and Mariposa Bypass along the San Joaquin River	7
3	Measuring understory cover along transect	8
4	Measuring overstory cover along transect	9
5	San Joaquin River discharge (cfs) measured at USGS gage 11251000 below Friant, California for Water Years 2011 to 2013	11

Contents, continued

Figures	<i>Page</i>
6 Relative percent cover of lifeforms in the understory layer of vegetation transects in the upper (A) and lower (B) reaches of the San Joaquin River from 2011 to 2013	13
7 Cluster analysis using Bray-Curtis similarities of plant species based on square root transformation of total percent cover data between reaches along the San Joaquin River in years 2011 to 2013.....	19
8 Locations of piezometers 7 and 8 and vegetation transects 1 and 2 within Reach 2B	21
9 Depth to groundwater at wells 7 and 8 and San Joaquin River discharge at gauges SJN and SJB from March to December 2013 in Reach 2B.....	21
10 Delta button celery (<i>Eryngium racemosum</i>), a State-listed endangered plant, was detected in transects within the Merced NWR in the East Side Bypass Reach, June 2012	26

Introduction

Background

In 2006, the Department of the Interior entered into the San Joaquin River Settlement (Settlement) in *NRDC et al., v. Kirk Rodgers et al.* The Settlement was subsequently approved by the Court in October 2006 and the San Joaquin River Restoration Settlement Act (Act), Public Law 111-11, authorizes and directs the Secretary of the Interior to implement the Settlement. The San Joaquin River Restoration Program (SJRRP) is a comprehensive long-term effort to restore flows and a self-sustaining Chinook salmon fishery to the San Joaquin River from Friant Dam to the confluence of Merced River, while reducing or avoiding adverse water supply impacts.

Historically, riparian vegetation in California's Central Valley was typical of a dynamic system largely driven by annual flooding and a long summer drought (Thompson 1961 as cited in Stillwater Sciences 2003a). Vegetation recruitment and survival were maintained through annual flooding via floodplain inundation, scour, and sediment deposition. Water availability during summer drought was the primary factor structuring vegetation establishment and distribution. This cycle of flooding and drought was – and still is – important to pioneer woody plant species, primarily willows (*Salix* spp.) and cottonwoods (*Populus* spp.), which rely on floods for bare seed beds, water, and nutrients, and which grow roots quickly to reach permanent water tables and a secure bank footing to resist subsequent floods (Braatne et al. 1996 as cited in Stillwater Sciences 2003a).

Riparian forests require periodic seedling recruitment and subsequent establishment to maintain the stand through time (Stillwater Sciences 2003a). A mature riparian zone typically consists of a mosaic of vegetation types of various ages and species. Commonly, mixed riparian forests occupy mid-elevation floodplain sites, and valley oak woodland and savannah occupy the oldest and driest floodplain sites such as high terraces and cut banks. Riparian vegetation dynamics are tightly coupled with river processes. Along geomorphically active streams, cottonwoods and willows are typically among the first species to colonize bare stream banks and bars. These species, with traits such as high seed output and rapid growth rates, tend to establish in bands parallel to the channel, with the youngest stands occurring closest to the active channel (Gregory et al. 1991, McBride and Strahan 1984, Walker and Chapin 1986 as cited in Stillwater Sciences 2003a). Each band of vegetation represents a separate recruitment event. Over time, pioneer vegetation traps sediment and adds litter and nutrient inputs to floodplain soils (Walker and Chapin 1986 as cited in Stillwater Sciences 2003a). As the floodplain develops and the riparian stand ages, changes in microclimate (shade, temperature, relative humidity) occur which often facilitates establishment of other riparian species such as Oregon ash (*Fraxinus latifolia*), box elder (*Acer negundo*), and valley oak (*Quercus lobata*). These “later successional” species typically produce larger seeds and are more shade-tolerant than the early pioneers, allowing them to persist in the seedbank

and germinate under the forest canopy when soil temperature and moisture conditions are adequate. Recruitment of these species is not as dependent on flow and sediment conditions as willows and cottonwoods.

Riparian vegetation along the San Joaquin between Friant Dam and the Merced River confluence has been significantly modified by agricultural development, hydrologic changes from operations of Friant Dam and the construction and operation of the flood control levees and bypass system. River regulation has created artificial hydrologic conditions, resulting in decreased peak flows, increased summer base flows, and reduction of physical processes such as scour and sediment deposition compared with historical conditions (Stillwater Sciences 2003a). Riparian pioneer tree populations that evolved with pre-regulation cycles of flooding and drought have decreased recruitment and altered topographic distributions relative to bank elevation and proximity to the channel (Strahan 1984, McBain and Trush 2000, Stillwater Sciences 2001 as cited in Stillwater Sciences 2003a). The reduction in riparian tree recruitment is compounded by human development on floodplains that has simultaneously removed over 90 percent of the historical riparian forests for fuel wood, agricultural and urban expansion, and floodplain mining (Katibah 1984 as cited in Stillwater Sciences 2003a). The San Joaquin River historically supported a much wider riparian corridor than is present under current conditions.

Reduced riparian vegetation along streambanks has decreased shaded riverine cover, organic inputs, water temperature control, and habitat structure (including inputs of large woody debris to aquatic habitats in the river), thus degrading aquatic habitat and fishery health. Important functions of the floodplain have also been reduced or eliminated, including flood flow retention and the ability for the channel to meander, which in turn increases both the risk of flooding and the amount of sediment deposited by flood flows.

In order to evaluate the establishment and development of riparian vegetation in response to Interim Flows, Reclamation's Technical Service Center (TSC) in Denver, CO and Mid-Pacific Region in Sacramento, CA established monitoring transects in river reaches 1A through 5 and including the East Side and Mariposa Bypasses. Monitoring began in August 2011 and will be conducted annually for comparison over time. In 2012, additional vegetation transects were established and monitored in river reach 5, which will also continue to be monitored annually. Hydrologic variables, including discharge and depth to groundwater as they relate to vegetation, were also incorporated in the monitoring program. Interim Flows were implemented in Water Year 2010; changes over the monitoring period beginning in 2011 will be evaluated.

Project Area

Vegetation transects were located in several reaches of the SJRRP Restoration Area (reach descriptions from CDWR 2002):

Reach 1A – River mile (RM) 267-243; Friant Dam to Highway 99 bridge at Herndon. This reach has the greatest diversity of vegetation types and the highest overall diversity

of plant species. It is also the most urbanized region of the project area, and has more gravel extraction and the least number of confining levees of any of the reaches. Riparian oak forest and mixed riparian forest are more commonly encountered in Reach 1A than downstream. Herbaceous and exotic vegetation types account for two-thirds (66.8 percent) of the total natural vegetation mapped, while approximately one-quarter (26.8 percent) is riparian forest. Woody scrub makes up less than seven percent (6.5 percent) of the total natural vegetation. The most common natural habitat types found here are: herbaceous (2701 acres), mixed riparian forest (526 acres), riparian oak forest (289 acres), willow scrub (290 acres), wetland/marsh (247 acres), and willow riparian (233 acres). The ratio of habitat per river mile is 194.2 acres/mi. (In these reach descriptions, CDWR (2002) describes “habitat” as naturally occurring vegetation, which excludes agricultural fields, open water, disturbed areas, or urban areas). In addition to woody exotic trees and giant reed (*Arundo donax*), scarlet wisteria (*Sesbania punicea*) is widespread in portions of Reach 1A. It has invaded wide areas of the floodplain in this and the subsequent Reach 1B, displacing willow scrub along the edge of the low-flow channel.

Reach 1B – RM 243-229; Highway 99 bridge to Gravelly Ford. This reach is more narrowly confined by levees than the upper section. The proportion of herbaceous and exotic vegetation is closer to one half of the total natural vegetation (55 percent), while the proportion of woody riparian vegetation is closer to one-third (30.6 percent) of the total and occurs mainly in narrow strips immediately adjacent to the river channel. Willow scrub is more abundant (14.3 percent) than in Reach 1A. Outside the levees and steep bluffs, the land use is nearly all agricultural. Scarlet wisteria was observed as far downstream as river mile 240. Giant reed patches are commonly encountered. The most abundant habitat types are herbaceous (300 acres) and mixed riparian (280 acres), followed by cottonwood riparian (193 acres), willow scrub (155 acres) and willow riparian (120 acres). This reach has the second lowest ratio of natural vegetation per mile—in 14 miles of channel, there is a little over one square mile of natural habitat (48 acres/mile).

Reach 2 (2A and 2B) – RM 229-205; Gravelly Ford to Mendota Pool. This reach is characterized by seasonal drying in the late summer and fall. The water table recedes into the porous substrate, creating a pronounced riparian drought nearly every year. There is about half as much riparian forest, proportionally, as in Reach 1 (15 percent of natural and naturalized vegetation), about the same proportion of woody scrub communities (13.5 percent) as Reach 1B, and more herbaceous vegetation (71 percent) than in Reach 1 overall. The most abundant habitat type by far is herbaceous (718.7 acres), followed by riparian scrub (302.8 acres), willow scrub (254.2 acres), riverwash (173.8 acres), willow riparian (165.4 acres), and cottonwood riparian forest (124.5 acres). The ratio of natural vegetation/river mile is 79.0 acres/mi., about 60 percent higher than in Reach 1B, but 40 percent of that in Reach 1A. Cultivated lands occupy nearly all the lands outside the river bottom. The character of the reach changes somewhat near Mendota Pool (RM 216-204). Downstream of the bifurcation structure at RM 216 (SW of which is found the large elderberry savanna), the riparian zone is very narrowly confined to a thin strip 3-10 meters wide bordering the channel. The herbaceous understory is however, very rich in

native species and a high proportion of the total vegetative cover is native plants, possibly due to the exclusion of cattle and other domestic stock from these thin habitat strips.

Reach 3 – RM 205-182; Mendota Pool to Sack Dam. The reach is characterized by a continuous flow within a very confined channel, seasonally low water (although not as dry as Reach 2), and narrow strips of riparian habitat along the river's edge. Adjacent lands are mostly under cultivation, although the city of Firebaugh borders the river's west edge for 3 miles. This reach has the smallest proportion of herbaceous habitat (25.2 percent) and the highest proportion of riparian forest (53.7 percent). Willow scrub occupies 21 percent of the total extent of natural vegetation. The most common habitats are cottonwood riparian (460.8 acres), willow scrub (230.5 acres), herbaceous (174.4 acres), and willow riparian (124.8 acres). Forty-seven and one-half acres of natural vegetation were mapped for every river mile in this reach, equivalent to the ratio found for Reach 1B.

Reach 4A – RM 182-148; Sack Dam to southern portions of the San Luis National Wildlife Refuge. This reach begins in cultivated and ends in public lands. Access for field verification and transects was denied in about half of this stretch. Reach 4A has the fewest habitat types and the lowest ratio of natural vegetation per river mile of any of the segments—only 502 acres of vegetation in this 34-mile segment (14.8 acres/mi.). The proportion of herbaceous habitats is typical of the San Joaquin River as a whole—about two-thirds (67.7 percent), while the proportion of forest is 22.4 percent and the proportion of woody scrub is 5 percent. The most common habitats are herbaceous (177.2 acres), willow riparian forest (89.1 acres), riverwash (65.2 acres), and riparian scrub (56.7 acres).

Reach 4B – RM 136 to 148; continues through public lands to the confluence with Bear Creek. Cultivated fields border approximately nine miles of the river's eastern bank. The floodplain is broad between widely spaced levees and the water table is nearer the surface than in some of the other reaches. These factors, along with a much lower level of disturbance to the native landscape on the public lands, create vast areas of natural habitat, compared to the upstream reaches. The ratio of natural habitat per river mile increases thirty-five-fold over that of Reach 4A, with a similar ratio continuing to the Merced River confluence (512.8 acres/mi. in Reach 4B). The most common vegetation type by far in this reach is herbaceous vegetation (4175 acres), followed by willow riparian forest (701.2 acres), wetland/marsh (377.7 acres), and willow scrub (132.1 acres). Giant reed was not seen in this reach.

Reach 5 – RM 118 to 136; confluence with Bear Creek to the confluence with the Merced River. Eight miles of this reach are adjacent to cultivated lands on the eastern bank, while the rest is bordered by relatively undisturbed natural habitat of private duck clubs and State and federal lands designated as refuges and parks. The natural habitat mapped per mile is similar to that of Reach 4B: 508 acres/mi. The characteristic habitat type of this reach is herbaceous vegetation, with 7,239 acres spreading over the wide floodplains of the San Luis Wildlife Refuge and the North Grasslands Wildlife Area. Following in predominance are willow riparian (972.6 acres), wetland/marsh (532.02

acres), and willow scrub (86 acres). The amount of wetlands encountered in the 30 river miles of Reach 4B and Reach 5 total more than twice that contained in the 119 miles of Reaches 1 through 4A.

Methods

Vegetation Transects

In 2011, twenty permanent vegetation transects were established within river reaches 1A, 1B, 2A, 2B, 3, 4A, 4B2 (*i.e.* San Luis National Wildlife Refuge (NWR)), and the East Side and Mariposa Bypasses (Figures 1 and 2). In 2012, two permanent vegetation transects were established within river reach 5 (Figure 2). Due to the large project area (over 150 RM), it was feasible to locate and monitor two transects within each reach with the exception of the East Side Bypass, where four transects were placed, two of which were in the Merced NWR. Transects were placed in areas adjacent to the river channel within the active floodplain. These sites are subsequently subject to seasonal changes in water and nutrient input and scour and sediment deposition. These transects are not representative of vegetation types across entire reaches, but were chosen to identify potential vegetation change over time resulting from Interim Flows. Aerial photos of vegetation transects by river reach are shown in Figures A-1 to A-11 in Appendix A.

Plant cover, composition, and overstory height and stem density were collected along each transect. Habitat variable ratings were determined for the area encompassing the transect. The length of each transect was determined by the extent of the floodplain and varied from 35 to 100 meters (m). Waypoints for each end of transects are listed in Appendix B. Forms used to collect data are included in Appendix C.

Timing

Monitoring will be conducted annually during spring or summer months depending on flow levels with the objective of collecting data at similar river phases and when vegetation is at comparable stages of development each year.

Understory Vegetation

For understory measurements, cover and species composition were measured either every 0.5 or 1 m along the transect depending on the length of the transect. The point-intercept method was used, which entailed recording the first “hit” for herbaceous plants and for woody species under 1 m tall by species (Figure 3). If a plant was not intercepted, then bare soil, litter, rock, or water were recorded. The location and extent of invasive weed species were documented when encountered.




Figure 1.—Location of upstream vegetation transects in Reaches 1, 2, and 3 along the San Joaquin River.




Figure 2.—Location of downstream vegetation transects in Reaches 4 and 5, East Side Bypass (ESB), and Mariposa Bypass (MB) along the San Joaquin River.



Figure 3—Measuring understory cover along transect.

Overstory Vegetation

The line-intercept method was used for measuring woody overstory cover. Overstory cover was measured along the transect by noting the point along the tape where the canopy began and the point at which it ended for each woody species over 1 m tall (Figure 4). Because species overlapped in some cases, the sum of the cover for all species did not necessarily reflect the actual percentage of overstory cover along the tape. The percentage of the tape covered by overstory was also calculated. The height of the tallest vegetation within each continuous stretch of the same species was measured.

Stem Density

Woody stem density was determined by using a meter stick to measure one meter outward on the upstream side of the transect. All woody stems within the one meter belt transect were counted and recorded by size into 4 classes by species (see Figure C-3 in Appendix C for descriptions of size classes).

Habitat Variables

A riparian systems model (Stein et al. 2000) was used to rank riparian condition. This qualitative model (riparian rank) includes spatial and structural diversity of native woody plants, contiguity of dominant vegetation, invasive vegetation, hydrology, topographic complexity, characteristics of flood-prone areas, and biogeochemical processing. These criteria consider the interaction between geology, hydrology, and organic and inorganic inputs to the system. Each criterion is scored between 0 and 1.0 and scores are added so that the “best” rank is an 8. See Figure C-4 in Appendix C for a listing of the variables and descriptions.



Figure 4—Measuring overstory cover along transect.

Statistical Analysis

Total cover and density data were compared between sampling periods for upstream reaches (Reaches 1 to 3) and downstream reaches (Reaches 4, 5, and Bypasses) to evaluate any statistically significant changes in vegetation over time. The repeated measures ANOVA was applied to test for relationships between cover or density and year, while Fisher's least significant difference (LSD) procedure was used as a multiple comparison test to evaluate statistically significant differences between years ($\alpha=0.05$) utilizing StatGraphics statistical software. The Fisher's LSD analysis is a post-test to the repeated measures ANOVA and provides a more focused analysis of individual years. Primer (Plymouth Routines in Multivariate Ecological Research; see www.primer-e.com) statistical software was used to create a similarity matrix and cluster diagram of plant species between reaches and between years using the Bray-Curtis measure.

Photo Stations

Two digital photographs were taken at each end of the transect – one toward the transect and one facing outward. These photos will provide visual documentation of vegetation height, density, species composition, and general site development for comparison over time.

Groundwater Monitoring

Reclamation installed piezometers to measure groundwater levels on vegetation transects in Reach 2B and 4A following the 2012 vegetation monitoring season. Groundwater recession rates have been closely tied to riparian vegetation establishment and survival in the San Joaquin Valley and elsewhere (Stillwater Sciences 2003b). Groundwater data from wells installed near transects will be used to determine if any correlations are observed between growth and development of vegetation and water table levels.

Results

Vegetation Transects

See Appendix D for a plant list of all herbaceous and woody species detected in transects within all reaches over 3 years of monitoring.

Timing

Vegetation transects were monitored June 11-14, 2012 and June 24-27, 2013 which was approximately 5-7 weeks earlier than in 2011, when monitoring was conducted August 1-4. 2011 was a wet year and Friant Dam was in flood operations through mid-July. Monitoring was not feasible until relatively late in the summer due to high river levels and inundated sites.

The extreme differences in river discharges between years can be seen in Figure 5, which shows the hydrograph for Water Year 2011 to 2013 along the San Joaquin River approximately 3.7 kilometers (2.3 miles) downstream of Friant Dam. River levels were similar in August of 2011 and June of 2012 and 2013.

Understory Vegetation

Seventy-nine annual and perennial species were identified while measuring understory vegetation along transects in all river reaches combined over 3 years of monitoring. The average total percent cover by individual species, life-form (*i.e.* native or introduced shrubs < 1m, grasses, and forbs) and cover type (*i.e.* plant, litter, bare ground, rock, water) found in the understory layer are shown for upstream Reaches 1A, 1B, 2A, 2B, and 3 in Table E-1, Appendix E and for downstream Reaches 4A, ESB, MB, 4B2, and 5 in Table E-2, Appendix E. A summary of total percent cover in the understory layer by cover type is shown in Table 1.

General trends in total cover among upstream reaches were an increase in total and introduced species cover while litter either changed little or increased. The exception was Reach 3, in which introduced species also increased but total cover decreased. Among downstream reaches, the general trend was a decrease in total cover, not much change in




Figure 5.—San Joaquin River discharge (cfs) measured at USGS gage 11251000 below Friant, California for Water Years 2011 to 2013. Source: United States Geological Survey.

Table 1.—Average total percent cover by type in the understory layer of vegetation transects along the San Joaquin River from 2011 to 2013.

Average Total Percent Understory Cover - Upstream Reaches															
Species	River Reach														
	1A			1B			2A			2B			3		
	2011	2012	2013	2011	2012	2013	2011	2012	2013	2011	2012	2013	2011	2012	2013
Native Plant Cover	14.5	7.5	15.5	9.5	4.0	10.5	9.0	2.5	2.5	19.1	6.3	14.8	60.4	13.6	17.7
Introduced Plant Cover	8.5	41.0	40.0	2.5	16.5	16.5	10.0	17.5	24.0	6.7	19.3	14.2	13.0	17.9	43.1
Total Plant Cover	23.0	48.5	55.5	12.0	20.5	27.0	19.0	20.0	26.5	25.8	25.6	29.0	73.4	31.5	60.8
Litter	33.5	39.0	39.0	20.0	40.5	42.5	16.5	17.5	15.5	22.9	32.8	33.3	16.0	56.9	34.5
Bare	37.0	6.0	2.5	55.0	32.5	30.5	45.0	43.0	58.0	51.3	41.6	37.8	10.6	11.6	4.8
Rock	6.0	5.0	2.5	13.0	6.5	0.0	19.5	19.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Water	0.5	1.5	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Total Cover	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Average Total Percent Understory Cover - Downstream Reaches														
Species	River Reach													
	4A			ESB			MB			4B2			5	
	2011	2012	2013	2011	2012	2013	2011	2012	2013	2011	2012	2013	2012	2013
Native Plant Cover	11.3	6.6	no data	29.8	25.2	21.5	63.5	17.0	25.0	41.5	28.8	40.3	19.5	11.0
Introduced Plant Cover	7.0	14.0	no data	17.5	18.5	15.5	30.5	26.5	26.5	39.9	37.9	31.4	26.0	34.0
Total Plant Cover	18.3	20.6	no data	47.3	43.7	37.0	94.0	43.5	51.5	81.4	66.7	71.7	45.5	45.0
Litter	7.5	26.7	no data	22.7	39.5	49.3	4.5	55.0	44.0	11.8	32.8	28.4	42.0	49.0
Bare	74.2	52.7	no data	30.0	16.8	13.8	1.5	1.5	4.5	6.8	0.5	0.0	12.5	6.0
Total Cover	100.0	100.0	no data	100.0	100.0	100.1	100.0	100.0	100.0	100.0	100.0	100.1	100.0	100.0

the percentage of introduced species, and an increase in litter. The only exception was in Reach 5, where total cover stayed the same. There were not drastic changes in total cover of native species among reaches except in Reach 3 and the Mariposa Bypass, where there were relatively large decreases in native species cover compared to 2011.

Species richness – or the number of species detected along the transect incorporating all measurement methods – was highest in the East Side Bypass in 2013, where 20 different herbaceous plant species were detected (Table 2). Species richness was consistently among the highest in the East Side Bypass and Reach 4B2 over the 3 years, with most species introduced. Reach 5 had the highest herbaceous species richness in 2012, but the number of species detected dropped from 19 to 12 in 2013; the majority of species detected in this reach were native. Native plant species richness decreased over the monitoring period in most reaches.

Table 2.—Species richness (number of species detected) for herbaceous and woody plants by reach along the San Joaquin River from 2011 to 2013.

Reach	Number of herbaceous species detected								
	Total			Native			Introduced		
	2011	2012	2013	2011	2012	2013	2011	2012	2013
1A	13	12	14	8	6	7	5	6	7
1B	3	6	7*	1	3	3	2	3	3
2A	15	10	8	7	4	2	8	6	6
2B	8	10	12	5	4	7	3	6	5
3	14	4	11	10	1	4	4	3	8
4A	9	7	na	4	3	na	5	4	na
ESB	16	16*	20*	9	7	8	7	6	11
MB	14	10*	11	8	4	6	6	5	5
4B2	16	16	16	8	6	7	8	10	9
5	na	19*	12	na	8	8	na	7	4
Reach	Number of woody species detected								
	Total			Native			Introduced		
	2011	2012	2013	2011	2012	2013	2011	2012	2013
1A	6	6	6	6	6	6	0	0	0
1B	6	6	7	4	4	5	2	2	2
2A	2	3	3	1	2	2	1	1	1
2B	3	4	4	3	4	4	0	0	0
3	2	2	2	2	2	2	0	0	0
4A	1	1	na	1	1	na	0	0	na
ESB	1	0	0	1	0	0	0	0	0
MB	no woody species								
4B2	1	1	1	1	1	1	0	0	0
5	na	1	1	na	1	1	na	0	0

*Includes unidentified plants; native/introduced status not determined

Relative percent understory cover of lifeforms by reach is shown in Figure 6. In 2011, native forbs were generally the most common lifeforms detected. Dominant lifeforms shifted to primarily introduced grasses in the upper reaches in 2012 and 2013. In the lower reaches, introduced forbs comprised the largest proportion of lifeforms




Figure 6.—Relative percent understory cover of lifeforms in vegetation transects by reach of the San Joaquin River from 2011 to 2013. Data was not collected in Reach 4A in 2013.

documented in 2012, while in 2013 no lifeform dominated the species composition in these reaches. Native species were dominant relative to introduced species among understory plants in all reaches in 2011; however this trend shifted in 2012, when relative cover of introduced species was higher than native species in all reaches with the exception of the East Side Bypass, where relative cover of native species was 58 percent (Table 3). This pattern was the same in 2013 except the proportion of native plant cover also increased in Reach 2B to 51 percent and in Reach 4B2 to 56 percent.

Table 3.— Proportion of native and introduced species in the understory and overstory layers by reach along the San Joaquin River from 2011 to 2013.

Relative Percent Cover												
Reach	Understory layer						Overstory layer					
	Native spp			Introduced spp			Native spp			Introduced spp		
	2011	2012	2013	2011	2012	2013	2011	2012	2013	2011	2012	2013
1A	63	16	28	37	84	72	100	100	100	0	0	0
1B	79	22	39	21	78	61	82	82	77	18	18	23
2A	53	13	9	47	87	91	100	100	100	0	0	0
2B	74	25	51	26	75	49	100	100	100	0	0	0
3	82	43	29	18	57	71	100	100	100	0	0	0
4A	62	32	NA	38	68	NA	no overstory		no data		no overstory	
ESB	51	58	58	49	42	42	100	no overstory		0	no overstory	
MB	73	39	49	27	61	51	no overstory					
4B2	68	43	56	32	57	44	100	100	100	0	0	0
5	NA	43	24	NA	57	76	NA	100	100	100	NA	0

Overstory Vegetation

There were 12 woody species detected in the overstory layer (woody species > 1m in height) of all transects combined (Table 4). No overstory was recorded along transects within Reach 4A and the Mariposa Bypass in any year. There was little change in total overstory cover from 2011 to 2013 in any reach except 2B, where cover increased from 7.2 to 18.4 percent (Table 4). In 2013, total percent overstory cover was highest in the most downstream reach 5 (70.8 percent) followed by the uppermost reaches 1A and 1B, with estimates of 44.5 and 58.2 percent, respectively. The average height of the tallest overstory shrubs/trees within each stretch by species is also shown in Table 4.

In general, woody species richness was directly related to proximity to Friant Dam, with only upstream Reaches 1A, 1B, and 2B having more than 2 woody species in the plant composition.

The vast majority of overstory trees and shrubs were comprised of native species relative to introduced (Table 3). The only overstory introduced species recorded were giant reed (technically a grass but also categorized as a shrub; USDA - NRCS 2012) and scarlet wisteria. Both were documented in overstory measurements in Reach 1B; scarlet wisteria was also noted in density measurements in Reach 2A. Both species are classified as a noxious weed in California.

Stem Density

Density of woody plants by size class and species is listed in Table 5. No stems were detected in the one meter belt associated with transects in Reach 4B2 and the East Side and Mariposa Bypasses. Highest densities were found in upstream Reaches 1A, 1B, 2B, and 5 in all years. Of these, Reach 1B had the highest density in 2013 (4.68 stems / m²), after a decrease in 2012. The relatively high number of stems in 2013 were mostly due to beaver browse of Goodding's willow that resulted in numerous resprouts. Within this reach, most stems were willow of size classes 2 and 3. Reach 1A also had a relatively large increase in density in 2013 which could be attributed to a high number of oak

Table 4.— Total percent cover and average height of woody overstory species (>1 m) detected in vegetation transects for upstream reaches (A) and downstream reaches (B) of the San Joaquin River from 2011 to 2013. No overstory species were documented in transects within Reaches 4A and Mariposa Bypass in any year.

Reach	Year	Native species																				Introduced species					Total canopy*	
		White alder		Button bush		Oregon ash		Fremont cottonwood		Valley oak		Sandbar willow		Gooding's willow		Red willow		Arroyo willow		Black elderberry		Total native	Giant reed		Scarlet wisteria			Total intro
		Tot % cov	Avg. Ht. (m)	Tot % cov	Avg. Ht. (m)	Tot % cov	Avg. Ht. (m)	Tot % cov	Avg. Ht. (m)	Tot % cov	Avg. Ht. (m)	Tot % cov	Avg. Ht. (m)	Tot % cov	Avg. Ht. (m)	Tot % cov	Avg. Ht. (m)	Tot % cov	Avg. Ht. (m)	Tot % cov	Avg. Ht. (m)	Tot % cov	Tot % cov	Tot % cov	Avg. Ht. (m)	Tot % cov		Avg. Ht. (m)
1A	2011	0.6	4.3	7.1	2.2	8.9	10.3	0		21.1	12.1	5.7	2.2	0		0		4.7	4.9	0		48.0	0		0		0.0	45.2
	2012	0		9.0	3.4	17.6	15.0	0		20.0	21.3	9.4	2.4	0		0		5.5	3.8	0		61.5	0		0		0.0	46.7
	2013	2.0	4.3	5.3	2.9	16.3	10.1	0		21.7	14.6	6.3	2.7	0		0		8.6	4.1	0		60.2	0		0		0.0	44.5
1B	2011	0		7.6	2.4	0		4.1	2	0		22.1	2.8	11.8	3.5	0		0		0		45.5	9.8	4.4	0.3	1.3	10.1	54.9
	2012	0		7.8	2.7	0		2.7	3.7	0		28.3	2.8	12.7	3.5	0		0		0		51.5	9.5	5.2	1.6	1.7	11.1	58.9
	2013	0		8.8	3.2	0		2.9	4.1	0		21.0	2.4	12.8	2.8	0		0		0		45.5	9.1	5.1	4.4	1.8	13.5	58.2
2A	2011	0		0		0		0		0		0		0		1.1	3.9	0		0		1.1	0		0		0.0	1.1
	2012	0		0		0		0		0		0		0		0.9	4.1	0		0		0.9	0		0		0.0	0.9
	2013	0		0		0		0		0		0		0		3.8	2.9	0		0		3.8	0		0		0.0	3.8
2B	2011	0		0		0		0		0		0		0		4.4	5.9	0		2.8	4.2	7.2	0		0		0.0	7.2
	2012	0		0		0		0		0		0		0		12.5	6.0	0		6.9	4.8	19.4	0		0		0.0	19.4
	2013	0		0		0		0.2		0		0.7		0		10.3	3.0	0		7.3	5.7	18.5	0		0		0.0	18.4
3	2011	0		0		0		16.7	15.0	0		0		0.8	3.0	0		0		0		17.4	0		0		0.0	17.4
	2012	0		0		0		14.4	15.0	0		0		2.8	3.0	0		0		0		17.2	0		0		0.0	16.1
	2013	0		0		0		13.7	23.0	0		0		7.2	3.6	0		0		0		20.9	0		0		0.0	18.3
ESB	2011	0		0		0		0		0		0		0.2	na	0		0		0		0.2	0		0		0.0	0.2
	2012	0		0		0		0		0		0		0		0		0		0		0.0	0		0		0.0	0.0
	2013	0		0		0		0		0		0		0		0		0		0		0.0	0		0		0.0	0.0
4B2	2011	0		0		0		0		0		0		9.3	8.6	0		0		0		9.3	0		0		0.0	9.3
	2012	0		0		0		0		0		0		8.5	10.5	0		0		0		8.5	0		0		0.0	8.5
	2013	0		0		0		0		0		0		9.5	10.2	0		0		0		9.5	0		0		0.0	9.5
5	2012	0		0		0		0		0		0		62.8	4.4	0		0		0		62.8	0		0		0.0	62.8
	2013	0		0		0		0		0		0		70.8	6	0		0		0		70.8	0		0		0.0	70.8

*Total canopy may not equal sum of all species due to overlap

Table 5.—Density of woody plant species by size class and species in river Reaches 1A to 2B along the San Joaquin River from 2011 to 2013.

Average # stems/m ²													
Species	Size class*	Reach											
		1A			1B			2A			2B		
		2011	2012	2013	2011	2012	2013	2011	2012	2013	2011	2012	2013
Giant reed	1	0	0	0	0	0	0	0	0	0	0	0	0
	2	0	0	0	0	0	0	0	0	0	0	0	0
	3	0	0	0	0.22	0.12	0.07	0	0	0	0	0	0
	4	0	0	0	0	0	0	0	0	0	0	0	0
Button bush	1	0.01	0	0	0.01	0	0	0	0	0	0	0	0
	2	0.07	0.14	0.01	0.22	0.16	0.02	0	0	0	0	0	0
	3	0.18	0.15	0.05	0.28	0.12	0.25	0	0	0	0	0	0
	4	0.02	0	0	0	0	0	0	0	0	0	0	0
Oregon ash	1	0.04	0	0	0	0	0	0	0	0	0	0	0
	2	0	0	0.05	0	0	0	0	0	0	0	0	0
	3	0.01	0	0.04	0	0	0	0	0	0	0	0	0
	4	0.01	0.02	0.01	0	0	0	0	0	0	0	0	0
Fremont cottonwood	1	0	0	0	0	0	0	0	0	0	0	0	0
	2	0	0	0	0	0	0	0	0	0	0	0.07	0.02
	3	0	0	0	0	0	0	0	0	0	0	0	0
	4	0	0	0	0	0	0	0	0	0	0	0	0
Valley oak	1	0.01	0	1.79	0	0	0	0	0	0	0	0	0
	2	0	0	0	0	0	0	0	0	0	0	0	0
	3	0.01	0.01	0.01	0	0	0	0	0	0	0	0	0
	4	0.01	0.01	0.01	0	0	0	0	0	0	0	0	0
Sandbar willow	1	0.01	0.04	0	0	0	0	0.19	0	0	0.01	0.02	0.01
	2	0.14	0.02	0.05	2.49	0.84	0.35	0.04	0.04	0.01	0	0.11	0.06
	3	0.31	0.61	0.28	1.14	0.95	1.75	0	0	0	0	0.05	0.17
	4	0.02	0	0	0.01	0	0	0	0	0	0	0	0
Goodding's willow	1	0	0	0	0	0	0	0	0	0	0	0	0
	2	0	0	0	0	0	2.04	0	0	0	0	0	0
	3	0	0	0	0.05	0.16	0.17	0	0	0	0	0	0
	4	0	0	0	0	0	0	0	0	0	0	0	0
Red willow	1	0	0	0	0	0	0	0	0.13	0	1.56	0.80	0
	2	0	0	0	0	0	0	0	0.12	0.03	0	1.78	0.38
	3	0	0	0	0	0	0	0	0	0	0.06	0.68	0.46
	4	0	0	0	0	0	0	0	0	0	0	0	0
Shining willow	1	0.04	0	0	0	0	0	0	0	0	0	0	0
	2	0.22	0.05	0.36	0	0	0	0	0	0	0	0	0
	3	0	0.10	0.03	0	0	0	0	0	0	0	0	0
	4	0	0	0	0	0	0	0	0	0	0	0	0
Black elderberry	1	0	0	0	0	0	0	0	0	0	0	0	0
	2	0	0	0	0	0	0	0	0	0	0	0	0
	3	0	0	0	0	0	0	0	0	0	0.32	0.10	0.21
	4	0	0	0	0	0	0	0	0	0	0	0	0
Scarlet wisteria	1	0	0	0	0	0	0	0.02	0.02	0.02	0	0	0
	2	0	0	0	0.02	0.01	0	0	0	0	0	0	0
	3	0	0	0	0	0	0.03	0	0	0	0	0	0
	4	0	0	0	0	0	0	0	0	0	0	0	0
Total by size class	1	0.10	0.04	1.79	0.01	0	0	0.21	0.15	0.02	1.57	0.82	0
	2	0.43	0.20	0.47	2.73	1.02	2.41	0.04	0.16	0.04	0	1.96	0.47
	3	0.50	0.87	0.41	1.69	1.35	2.27	0	0	0	0.38	0.83	0.84
	4	0.06	0.03	0.02	0.01	0	0	0	0	0	0	0	0
TOTAL stems/m ²		1.08	1.13	2.68	4.44	2.37	4.68	0.25	0.31	0.06	1.95	3.62	1.32

*Size classes: 1= current year's seedling
2= <1 m in ht
3= >1 m in ht and <10 cm DBH
4= >10 cm DBH

Table 5, cont'd.—Density of woody plant species by size class and species in river Reaches 3, 4A, and 5 along the San Joaquin River from 2011 to 2013.

Average # stems/m ²								
Species	Size class*	Reach						
		3			4A**		5	
		2011	2012	2013	2011	2012	2012	2013
Fremont cottonwood	1	0	0	0	0	0	0	0
	2	0.07	0	0	0	0	0	0
	3	0	0	0	0	0	0	0
	4	0	0	0	0	0	0	0
Goodding's willow	1	0	0	0	0.02	0	0	0
	2	0	0	0	0.41	0.86	0.76	0
	3	0.04	0.04	0.01	0	0	0.58	0.72
	4	0	0	0	0	0	0	0.01
Total by size class	1	0	0	0	0.02	0	0	0
	2	0.07	0	0	0.41	0.86	0.76	0
	3	0.04	0.04	0.01	0	0	0.58	0.72
	4	0	0	0	0	0	0	0.01
TOTAL stems/m ²		0.11	0.04	0.01	0.43	0.86	1.34	0.73

**4A was not sampled in 2013

seedlings (size class 1) detected along the transect. Densities in Reach 2B decreased somewhat substantially in 2013 from 3.62 to 1.32 stems/m² (mostly reductions in the number of red willow stems), as did densities in Reach 5, decreasing from 1.34 to 0.73 Goodding's willow stems/m².

Habitat Variables

The highest ranking habitat variables (>7.0) were found in the uppermost reaches 1A and 1B in 2013 (Table 6). These sites were rated relatively close to the highest possible ranking of 8.0, which indicates excellent riparian condition. Reaches with a moderate ranking (between 5.0 and 7.0) were 2A, 2B, and 5. All other reaches ranked between 3.3 and 4.7. These sites typically ranked relatively low in variables "Coverage and Spatial Diversity", "Structural Diversity", "Micro- and Macrotopographic Complexity", and "Biogeochemical Processes". Changes in habitat variables from 2011 to 2013 differed among reaches with the greatest differences in Reach 2B (increase of 1.45) and Reach 3 (decrease of 1.1).

Statistical Analysis

Statistical comparisons between years were performed on upstream reaches (Reaches 1A to 3) and downstream reaches (Reaches 4A to 5 and Bypasses) separately. This division of areas was based on cluster analysis of species composition similarities between reaches (Figure 7). There is a general division between the upper reaches (to the left on the graph in Figure 7) and lower reaches (to the right).

Statistical differences in the upstream reaches were in total cover of litter (significantly less in 2011 than 2012) and in total cover of introduced understory species (significantly less in 2011 than in both 2012 and 2013; Table 7). The only statistical difference identified in downstream reaches was significantly less litter cover in 2011 than in other years.

Table 6.—Ranking of habitat variables as an indicator of riparian condition by reach along the San Joaquin River from 2011 to 2013.

Reach	Year	Coverage & Spatial Diversity	Structural Diversity	Contiguity of Habitats	% Invasive Woody Vegetation	Hydrology	Micro- & Macrotopographic Complexity	Characteristics of Flood-prone Area	Biogeochemical Processes	Total Score*
1A	2011	1.00	0.70	1.00	1.00	1.00	1.00	1.00	0.90	7.60
	2012	0.90	0.65	1.00	1.00	1.00	0.80	1.00	0.90	7.25
	2013	0.95	0.75	0.95	1.00	1.00	0.90	1.00	0.90	7.45
1B	2011	1.00	0.75	0.80	0.90	1.00	0.85	1.00	0.80	7.10
	2012	1.00	0.85	1.00	0.90	1.00	0.90	1.00	0.95	7.60
	2013	1.00	0.80	0.90	1.00	1.00	1.00	1.00	0.95	7.65
2A	2011	0.25	0.25	0.80	1.00	1.00	0.80	0.90	0.35	5.35
	2012	0.40	0.40	0.90	1.00	1.00	0.50	1.00	0.30	5.50
	2013	0.35	0.40	0.80	1.00	1.00	0.95	1.00	0.40	5.90
2B	2011	0.25	0.40	0.90	1.00	1.00	0.35	0.90	0.30	5.10
	2012	0.40	0.40	0.90	1.00	1.00	0.70	0.90	0.50	5.80
	2013	0.75	0.40	0.80	1.00	1.00	0.95	1.00	0.70	6.55
3	2011	0.45	0.60	0.65	1.00	0.80	0.70	0.80	0.60	5.60
	2012	0.55	0.50	0.80	1.00	0.90	0.25	0.70	0.50	5.20
	2013	0.40	0.40	0.80	1.00	0.60	0.35	0.45	0.50	4.50
4A**	2011	0.20	0.20	0.80	1.00	1.00	0.35	1.00	0.20	4.75
	2012	0.20	0.20	0.80	1.00	0.60	0.20	0.30	0.30	3.60
ESB	2011	0.18	0.18	0.90	1.00	0.50	0.28	0.83	0.40	4.25
	2012	0.23	0.25	0.90	1.00	0.30	0.28	0.33	0.28	3.55
	2013	0.25	0.18	0.80	1.00	0.30	0.20	0.40	0.30	4.53
MB	2011	0.20	0.25	1.00	1.00	0.30	0.20	0.30	0.40	3.65
	2012	0.25	0.30	0.80	1.00	0.60	0.20	0.30	0.35	3.80
	2013	0.20	0.20	0.80	1.00	0.45	0.20	0.25	0.20	3.30
4B2	2011	0.40	0.40	0.80	1.00	0.70	0.50	0.80	0.60	5.20
	2012	0.40	0.40	0.90	1.00	0.60	0.45	0.80	0.40	4.95
	2013	0.30	0.40	1.00	1.00	0.60	0.50	0.45	0.45	4.70
5	2012	0.50	0.70	1.00	1.00	1.00	0.90	1.00	1.00	7.10
	2013	0.40	0.80	1.00	1.00	0.95	0.90	0.80	1.00	6.85

* Possible score 0 (Poor) to 8 (Excellent)

**4A was not sampled in 2013




Figure 7.— Cluster analysis using Bray-Curtis similarities of plant species based on square root transformation of total percent cover data between reaches along the San Joaquin River in years 2011 to 2013 (Y11 to Y13). The most similar reaches are grouped together.

Table 7.— Statistical results comparing total plant cover and density over time for various parameters in upstream and downstream reaches of the San Joaquin River. Alpha = 0.05.

Upstream Reaches (1-3)			
Parameter	Repeated measures ANOVA		LSD
	P-value		Significant difference (P<0.05)
Total cover	Plant	P=0.466	No difference
	Litter	P=0.037	11<12
	Bare	P=0.350	No difference
	Native understory	P=0.060	No difference
	Introduced understory	P=0.015	11<12 and 13
	Overstory	P=0.936	No difference
	Salix spp	P=0.812	No difference
Density	Total	P=0.964	No difference
	Salix spp	P=0.996	No difference
Downstream Reaches (4-5 and Bypasses)			
Parameter	Repeated measures ANOVA		LSD
	P-value		Significant difference (P<0.05)
Total cover	Plant	P=0.335	No difference
	Litter	P<0.001	11<12 and 13
	Bare	P=0.096	No difference
	Native understory	P=0.091	No difference
	Introduced understory	P=0.966	No difference
	Overstory	P=0.488	No difference
	Salix spp	P=0.523	No difference
Density	Total	P=0.231	No difference
	Salix spp	P=0.231	No difference

Highlighted boxes = significant difference at the 95% confidence level

Photo Stations

Photographs taken from the end of vegetation transects the first three years of monitoring are shown in Appendix F. Differences in vegetation along transects within many of the reaches are evident when comparing the photos from 2011 (a very wet year) to photos from the other 2 years.

Groundwater Monitoring

Piezometers were installed in association with vegetation transects in Reach 2B in February 2013 (Figure 8). The hydrograph in Figure 9 shows groundwater depths at these wells from March through December and uses flow data gathered at Station SJB (approximately 1.5 mi upstream) and Station SJN (approximately 2 mi downstream). A correlation between flows and the depth of the water table is apparent, which indicates connectivity of the floodplain and river. In 2013, ground water within the floodplain (PZ7) remained at a shallow enough depth (<4 ft) necessary to sustain woody riparian plant species.



Figure 8.—Locations of piezometers 7 and 8 and vegetation transects 1 and 2 within Reach 2B. Google Earth imagery August 2012.




Figure 9.—Depth to groundwater at wells 7 and 8 and San Joaquin River discharge at gauges SJN and SJB from March to December 2013 in Reach 2B.

Discussion

Following is a descriptive analysis comparing vegetation parameters from 2011 to 2013 by reach. The 2011 Water Year was very different hydrologically from 2012 and 2013 (Figure 4), which was likely a cause for many of the changes observed.

Reach 1A

This reach was the only one to show a perceptible increase in understory plant cover from 2011 to 2013, increasing from 23.0 to 55.5 percent in total cover (Table 1). This rise was due to a substantial increase in introduced species, from 8.5 to 40.0 percent; these species were exclusively introduced grasses (Table E-1, Appendix E). This change is evident in photos from 2011 to 2013 (Appendix F; Reach 1A, Transect 1; 1A-Toward transect and Transect 2; 2B-Toward transect). Low river banks at this site allow for overbank flooding, a condition that was prolonged in 2011 and likely deterred the establishment of introduced grasses, which are not as adaptable to inundation as native riparian species. In 2012 and 2013, conditions were drier and upland introduced grass species detected at the site (i.e. ripgut brome, softchess brome, foxtail chess, and rattail fescue) were able to thrive. The total bare cover decreased from 37.0 in 2011 to 2.5 percent in 2013, which was likely due to high flows and major flooding that caused scouring in 2011 and was not a factor in the other 2 years. The increase in understory introduced species resulted in a shift in dominance of native species at the site, with relative cover of understory native species decreasing from 63.0 percent in 2011 to 28.0 percent in 2013 (Table 3).

Understory species richness increased from 13 species detected to 14 over the course of monitoring, with the number of native species detected decreasing from 8 to 7 (Table 2). Total cover of overstory species within this reach remained essentially the same although the cover of native species increased from 48.0 percent in 2011 to 60.2 percent in 2013 (Table 4), indicating that although total canopy (i.e. the amount of overstory covering the transect) did not spread into open areas, native species canopy became more dense. This reach continued to have one of the highest diversity of woody species, with species richness remaining at 6 (Table 2). Stem density in Reach 1A increased from 1.08 to 2.68 stems/m² (Table 5). The large increase in 2013 was due to the detection of numerous oak seedlings. Recruitment of this species most likely indicates drier conditions since valley oak is more tolerant of drought and is less dependent on flow and sediment conditions than willows and cottonwoods. Also due to the high number of oak seedlings, the dominant size class shifted from Class 3 (>1 m in ht and <10 cm DBH) in 2011 and 2012 to Class 1 (current year's growth) in 2013. Finally, habitat variable rating decreased slightly from 7.6 to 7.45 (Table 6), although this reach still remained among the top in riparian health.

Reach 1B

This reach showed the second highest increase in total understory cover among reaches, rising from 12.0 percent in 2011 to 27.0 percent in 2013 (Table 1). There was a notable

increase in total percent litter (from 20.0 to 42.5 percent), while bare cover decreased from 55.0 to 30.5 percent. These results were presumably related to flows and flooding – high river discharge in 2011 caused scouring (higher bare cover) and lower flows in 2012 and 2013 led to the accumulation of litter. Total cover of introduced species increased from 2.5 to 16.5 percent, which was predominantly due to an increase in introduced grasses (Table E-1, Appendix E). A dominance of native herbaceous species in 2011 shifted to a dominance of introduced species in 2012 and 2013, with relative cover of native herbaceous species decreasing from 79 to 39 percent (Table 3). Species richness in Reach 1B increased from 3 herbaceous species detected in 2011 to 7 detected in 2013, and despite dominance of introduced species, native species richness increased from 1 to 3 (Table 2). Total overstory cover showed a negligible increase, with total cover of the introduced species scarlet wisteria increasing slightly from 0.3 to 4.4 percent from 2011 to 2013 (Table 4). Based on cover, the site was dominated by sandbar willow. Stem density remained the highest relative to other reaches in 2013 despite a notable decrease in 2012 from 4.44 to 2.32 stems/m² (Table 5). The increase in 2013 was associated with Goodding’s willow that had been browsed by beaver resulting in several resprouted stems. Habitat variable rating increased in Reach 1B, from 7.1 to 7.65 (Table 6), which ranked the highest among reaches in 2013. This rating was very near the highest possible score of 8, which indicates excellent riparian health.

Reach 2A

Total understory cover did not show major changes from 2011 to 2013, although native understory cover decreased from 9.0 to 2.5 (due to a drop in native forbs) and introduced herbaceous species cover increased from 10.0 to 24.0 percent (due to a rise in introduced grasses; Table 1). As a result, relative percent cover of native species decreased from 52.6 to 9.0 percent (Table 3). Herbaceous species richness dropped from 15 in 2011 to 8 in 2012, with native species richness decreasing from 7 to 2 (Table 2). Total overstory cover was only around 1.0 percent in 2011 and 2012, with only red willow detected in the overstory measurements (Table 4). In 2013, overstory cover increased to 3.8 percent due to the development of young willow in the lowermost portion of transects. Woody species richness, on the other hand, increased from 2 to 3, with stem density measurements increasing from 2 to 3 species counted. This was one of two reaches in which the noxious weed scarlet wisteria was documented. Despite the increase in woody cover in 2013, density decreased from 0.25 to 0.06 stems/m² (Table 5). Riparian condition was ranked moderately, increasing from 5.35 in 2011 to 5.9 in 2013 (Table 6), with lowest scores in the variables coverage and spatial diversity (i.e. cover and diversity of native riparian species), structural diversity (i.e. different size- and age- classes of riparian vegetation), and biogeochemical processes (i.e. vegetation with woody debris, leaf litter, detritus in channel).

Reach 2B

Similar to Reach 2A, total understory cover remained the same, with a slight decrease in total native cover (19.1 to 14.8 percent) and a slight increase in total introduced cover

(6.7 to 14.2 percent). Litter cover increased and bare cover decreased, both by around 10 percent (Table 1). Relative cover of native species decreased from 74 percent in 2011 to 25 percent in 2012 but increased to 51 in 2013 (Table 3, Figure 6), making native herbaceous species once again dominant at the site. Species richness increased from 8 herbaceous species detected to 10, although native species richness decreased from 5 to 4 (Table 2). This reach showed the largest increase in total overstory cover (7.2 to 18.4 percent; Table 4). There was also a large increase in stem density from 2011 to 2012 (1.95 to 3.62 stems/m²); however this number decreased to 1.32 stems/m² in 2013 (Table 5). The majority of stems shifted from size class 1 in 2011 to size class 2 in 2012 and to size class 3 in 2013. This increase in size would explain the decrease in overall stem density; as plants grow, their numbers decrease due to competition. Woody species richness increased from 3 to 4 (all native species). At this point in the study, Reach 2B shows the most potential for identifying change over time and effects from Interim Flows as a healthy riparian community appears to be developing. Young willow and cottonwood seedlings became established over the monitoring period, increasing in cover, size and richness (see comparison photos in Appendix F; Reach 2B, Transect 1; 1A). Piezometers installed at the site in 2013 indicate that hydrology is favorable for sustaining riparian vegetation, with a relatively shallow water table (< 4 ft) within the floodplain (Figure 9). Riparian condition remained moderate, increasing from 5.1 in 2011 to 6.55 in 2013 (Table 6), with lowest scores in the variables coverage and spatial diversity (i.e. cover and diversity of native riparian species), structural diversity (i.e. different size- and age- classes of riparian vegetation), and biogeochemical processes (i.e. vegetation with woody debris, leaf litter, detritus in channel).

Reach 3

This reach showed substantial changes over the study period. Total understory cover decreased from 73.4 percent in 2011 to 31.5 percent in 2012. Although total understory cover rebounded to 60.8 percent in 2013, this increase was correlated with a 30 percent rise in introduced species (Table 1). Reach 3A showed the largest decrease in total native cover among reaches over the course of monitoring, decreasing from 60.4 to 17.7 percent. Dominance in native herbaceous species relative to introduced species shifted in 2012; in 2013 native species composed 29 percent, down from 82 in 2011 (Table 3). This shift was related to an increase in introduced grasses (predominantly ripgut brome) at the site. Herbaceous species richness fell from 14 to 11 from 2011 to 2013 (Table 2). Native species richness experienced a relatively large drop, decreasing from 10 to 4. Total overstory cover remained the same (around 17 percent; Table 4), as did woody species richness (2 species detected; Table 2) from 2011 to 2013. Stem density decreased from 0.11 to 0.01, with stems detected only in size class 3 in 2012 and 2013 (Table 5). The difference in hydrologic conditions between the first and last 2 years of the study appeared to have large effects on herbaceous understory species within this reach, with drier conditions in 2012 and 2013 presumably causing cover, species richness and the native species component to decrease considerably. This site has high cutbanks, which prevented overbank flows – which occurred in 2011 – from occurring in 2012 and 2013. Reach 3 dropped from a moderate ranking to low in riparian condition (from 5.6 to

4.5) with a relatively large drop in the micro- and macrotopographic complexity (mixture of topographic features) and characteristics of flood-prone area variable rating.

Reach 4A

Due to access issues, this site was not monitored in 2013. From 2011 to 2012, total understory cover remained essentially the same at around 20 percent in Reach 4A, but consistent with most reaches over this period, native species cover decreased while introduced species cover increased (Table 1). Litter cover increased from 7.5 to 26.7 percent and bare cover decreased from 74.2 to 52.7, which was another common trend that was likely related to flooding and scouring in 2011 and low flows in 2012. Regardless, this site still had the highest percentage of bare ground in both 2011 and 2012. Native herbaceous species were no longer dominant relative to introduced species in 2012, decreasing from 61.9 to 31.8 relative percent cover (Table 3). No overstory cover was documented. Goodding's willow was, however, detected in both understory cover and density measurements in both years (woody species richness = 1; Table 2), indicating potential for willow riparian habitat if river conditions can sustain woody species along this reach. Stem density did increase slightly, from 0.44 to 0.86 stems/m²; all stems were detected in size class 2 in 2012, however, which suggested no recruitment of seedlings as in 2011, when stems were also detected in size class 1 (Table 5). Riparian condition was ranked relatively low, decreasing from 4.75 in 2011 to 3.6 in 2012, with low scores in most variables (Table 6).

East Side Bypass

Total understory cover within this reach remained between 37 and 47 percent in all years, with litter cover increasing from 22.7 to 49.3 percent and bare cover decreasing from 30.0 to 13.8 percent. Unlike other reaches, native herbaceous species remained dominant relative to introduced species throughout the entire monitoring period (Table 3), with native grasses the most common life form. Herbaceous species richness increased from 16 to 20, which is high relative to other transects, although fewer of the species detected were native (8 in 2013; Table 2). In 2012 and 2013, transects in this reach fell within exclusively herbaceous habitat; in 2011, 0.2 percent overstory cover was documented (Table 4) but no woody species were detected in understory or stem density measurements. The East Side Bypass received a score of 4.53 in riparian condition in 2013. Rankings were relatively low in all but 2 variables, which were contiguity of habitats and % invasive woody vegetation (only because there were no woody species). Two of four transects were located within the Merced NWR where an existing year-round water supply could make it difficult to identify changes in vegetation from Interim Flows. Delta button celery, which is a State-listed endangered plant, was detected in the Merced NWR within this reach (Figure 10). This plant was documented in transect monitoring in 2011 (Table E-2, Appendix E). Although it was not intercepted in transect cover measurements in 2012, it was observed in the area surrounding transect 3. It was not observed at all during 2013 sampling when the area showed signs of recent and heavy grazing.



Figure 10.—Delta button celery (*Eryngium racemosum*), a State-listed endangered plant, was detected in transects within the Merced NWR in the East Side Bypass Reach, June 2012.

Mariposa Bypass

Vegetation was strictly herbaceous with no woody species of any size detected in transects within the Mariposa Bypass in any year. The largest decrease in total understory cover from 2011 to 2013 occurred in this reach, dropping from 94.0 (highest total cover in 2011) to 51.5 percent (Table 1). Total cover of native species showed a substantial decrease from 63.5 percent in 2011 to 25.0 percent in 2013 while introduced species cover remained the same. The decrease in total plant cover was replaced with litter cover, which increased considerably, from 4.5 to 44.0 percent. Native species were no longer dominant in the understory layer, with relative percent cover decreasing from 73 to 49 percent from 2011 to 2013 (Table 3). Herbaceous species richness dropped from 14 in 2011 to 11 in 2013, with native species richness decreasing from 8 species detected to 6. Mariposa Bypass received the lowest ranking in riparian condition of all reaches studied in 2011 and 2013 with a habitat variable score of 3.30 in 2013. Improvement is needed in the variables coverage and spatial diversity (i.e. cover and diversity of native riparian species), structural diversity (i.e. different size- and age- classes of riparian vegetation), micro- and macrotopographic complexity (mixture of topographic features), and biogeochemical processes (i.e. vegetation with woody debris, leaf litter, detritus in channel).

Reach 4B2

Reach 4B2 had the highest total understory cover of all reaches in 2012 and 2013 at 66.7 and 71.7 percent, respectively, decreasing from 81.4 percent in 2011 (Table 1). There was an increase in litter cover over the study period, from 11.8 to 28.4 percent. Native species dominated in 2013 (as was so in 2011) with a relative cover of 56 percent (Table 3) and composed predominantly of native grasses (i.e. saltgrass; Table E-2 in Appendix E). Herbaceous species richness remained high at 16 in all years (Table 2), although less than half of species detected were native (from 8 in 2011 to 7 in 2013). While mature Goodding's willow was measured in total overstory cover all years (around 9 percent), no woody species were detected in understory cover or stem density measurements, indicating that recruitment is potentially low. Reach 4B2 received a moderately low ranking in riparian condition, slightly decreasing from 5.2 to 4.7. Lowest scores were given for the coverage and spatial diversity (i.e. cover and diversity of native riparian species), structural diversity (i.e. different size- and age- classes of riparian vegetation), and biogeochemical processes (i.e. vegetation with woody debris, leaf litter, detritus in channel) variables. This reach is located in the San Luis NWR and, like the Merced NWR in the East Side Bypass, has been supplied with year-round water; therefore hydrologic conditions may not change considerably and effects from Interim Flows may be difficult to determine.

Reach 5

Transects in this reach were first established and monitored in 2012; therefore only comparisons between 2012 and 2013 are discussed. Total understory remained the same, with a slight decrease in native cover and a slight increase in introduced species cover between years. The noxious weed Perennial pepperweed composed half of the understory species cover, increasing from 7.5 percent in 2012 to 22.5 percent in 2013. Although herbaceous species richness fell from 19 to 12, this reach had a high native species richness (8) in 2013 compared with other reaches. This site had the highest total overstory cover in 2013 at 70.8 percent, although Goodding's willow was the only woody species detected. Stem density decreased from 1.34 to 0.73 stems/m², composed of mostly Goodding's willow in size class 3. Riparian condition was rated relatively high at 7.1, following only the upper reaches 1A and 1B in ranking, indicating relatively good habitat quality in Reach 5.

Conclusions

Many of the vegetation parameters measured showed similar trends among reaches from 2011 to 2012, which was most likely a result of extreme differences in hydrologic conditions during the 2 water years. Exceptionally high flows (approximately 3,000 to 9,000 cfs) in January and April through July of 2011 created prolonged flooding (Figure 5). Water Years 2012 and 2013 were very similar to each other, with river discharge lower (approximately 1,000 cfs) throughout the year, prohibiting typical overbank flows.

Most changes in vegetation that were observed were likely due to differences in precipitation patterns and hydrology over the monitoring period, although some of the changes could probably be linked to Interim Flow implementation over the study period.

Total understory native cover decreased, as did herbaceous native species richness, in all reaches from 2011 to 2012. In 2013, most reaches showed increases in native cover and richness from 2012, but generally not to levels of 2011. Understory cover was composed of principally native species in 2011. This condition shifted to dominance of introduced species in the understory layer in 9 of 10 of reaches in 2012 and 6 of 9 reaches in 2013. Prolonged flooding in 2011 appeared to affect composition and cover of native understory species along the river and likely deterred the establishment of introduced species, which are generally not as adaptable to inundation as native riparian species. Native species were dominant in 2011, presumably because they are tolerant of anaerobic conditions and because of less competition from exotic species. The predominance of introduced species in 2012 and 2013 could be attributed to drier conditions. Understory cover was closely linked to discharge, with abundance of these shallow-rooted plant species dependent upon the amount of available water near the surface. Statistically, the percentage of introduced species was significantly greater in 2012 and 2013 than in 2011 in upper reaches (Reach 1A to 3).

Total litter cover increased in all reaches, which was a statistically significant change. In some cases, total bare cover decreased (however not statistically), which was likely due to high flows and major flooding that caused scouring in 2011 and was not a factor in 2012 or 2013.

Deeper rooted woody species in the overstory layer did not appear to be affected by lower flows, with total cover showing little change over the monitoring period. The exception was in Reach 2B, where overstory cover increased over the study period and where numerous willow seedlings were documented in 2011. Apparently these plants were established enough to withstand drier conditions in 2012 and 2013 and expand in cover. Groundwater monitoring has also shown that there is a relatively shallow water table associated with vegetation transects, which is conducive to successful restoration. Riparian recruitment sustained throughout monitoring at the Reach 2B site is likely related to implementation of Interim Flows. Willow recruitment was documented after the 2011 flood year and these new plants were maintained by Interim Flows and a relatively shallow water table (according to data collected in 2013).

Stem density was variable, increasing or decreasing among reaches, and no statistically significant changes were identified. A consistent trend, however, was a decrease in stem density in Size Class 1 (i.e. current year's growth), indicating little recruitment of new seedlings in 2012. This was most likely due to a lack of flooding in 2012 and 2013, a condition that is conducive to regeneration of willow and cottonwood species. The exception was in Reach 1A, where numerous valley oak seedlings were observed in 2013. This species is able to tolerate drier conditions than willow and cottonwood.

Generally, upstream reaches (*i.e.* 1A through 3, but particularly 1A and 1B) exhibited healthier riparian condition than downstream reaches, with greater cover, diversity, and density of woody species and higher habitat variable rankings. Subsequently, downstream reaches – with the exception of the wildlife refuges that sustain a year-round water supply – are likely to have a greater potential for showing effects from interim flows. Reach 5 (the furthestmost downstream reach in the study) was comparable to upstream reaches in that relatively high values were recorded for cover and density of woody species and habitat variable rankings. Reach 5 also had relatively high native herbaceous species richness. Because current riparian conditions at Reach 5 are relatively good, effects may be more difficult to detect here as well. Continued monitoring will determine if vegetative conditions have improved in transects along all reaches of the San Joaquin River included in this study.

Summary

The SJRRP Vegetation Monitoring Study evaluates the response of riparian vegetation to Interim Flows through comparison of transect data over time. Changes in vegetation may have implications for Friant Dam flow scheduling, habitat establishment supporting fish, and maintenance needs to convey flows. In 2011 SJRRP established transects, collected the first year of data, and ranked transects for riparian condition. This monitoring effort was continued in 2012 - with 2 new transects added to the study further downstream – and 2013. SJRRP will continue monitoring vegetation transects in 2014.

Literature Cited

- CDWR (California Department of Water Resources). 2002. Riparian vegetation of the San Joaquin River. Prepared by CDWR, San Joaquin District, Fresno for San Joaquin River Habitat Restoration Program, Fresno, California.
- Clark, K.R. and R.M. Warwick. 2001. Change in Marine Communities: An Approach to Statistical Analysis and Interpretation, 2nd edition. Plymouth, United Kingdom.
- Stein, E.D., Tabatabai, F., Ambrose, R.F., 2000. Wetland mitigation banking: a framework for crediting and debiting. *Environmental Management* 26:233-250.
- Stillwater Sciences. 2003a. Restoration Objectives for the San Joaquin River. Prepared by Stillwater Sciences, Berkeley, CA for Natural Resources Defense Council, San Francisco, CA and Friant Water Users Authority, Lindsay, CA.
- Stillwater Sciences. 2003b. Draft Restoration Strategies for the San Joaquin River. Prepared by Stillwater Sciences, Berkeley, CA for Natural Resources Defense Council, San Francisco, CA and Friant Water Users Authority, Lindsay, CA.
- USDA, NRCS. 2012. The PLANTS Database (<http://plants.usda.gov>, 19 January 2012). National Plant Data Team, Greensboro, NC 27401-4901 USA.

Appendix A

Aerial Photos of Vegetation Transects by River Reach
Upstream to Downstream



Reach 1A



Reach 1B




Reach 2A



Reach 2B



Reach 3



Reach 4A



East Side Bypass



East Side Bypass (Merced NWR), Transect 3



East Side Bypass (Merced NWR), Transect 4



Mariposa Bypass



Reach 4B2 (San Luis NWR)



Reach 5

Appendix B

Vegetation transect waypoints

All datum in NAD83.

Reach	Transect	Endpoint A		Endpoint B		Zone
		x	y	x	y	
R1A	1	255049	4091361	255081	4091315	11S
	2	254888	4091300	254940	4091218	11S
R1B	1	755779	4077621	755782	4077561	10S
	2	755580	4077600	755592	4077546	10S
R2A	1	751417	4074422	751327	4074469	10S
	2	751327	4074470	751230	4074504	10S
R2B	1	741586	4072746	741646	4072729	10S
	2	741552	4072759	741518	4072769	10S
R3	1	734778	4076749	734732	4076729	10S
	2	734713	4076882	734652	4076833	10S
R4A	1	718414	4100615	718463	4100664	10S
	2	718341	4100777	718393	4100780	10S
MB	1	703911	4119706	703910	4119656	10S
	2	703797	4119712	703795	4119662	10S
ESB	1	714230	4111882	714285	4111905	10S
	2	714194	4111872	714145	4111861	10S
	3	710325	4116027	710390	4116107	10S
	4	708217	4117404	708262	4117424	10S
R4B2	1	693717	4123312	693634	4123287	10S
	2	693670	4123484	693583	4123432	10S
R5	1	679685	4134377	679699	4134329	10S
	2	679658	4134367	679694	4134336	10S

Appendix C

Data collection forms

**San Joaquin Restoration Flows
Vegetation Monitoring
Tree Density
Site Characterization**

Date: _____

Transect: _____

Observers: _____

Density

Size class	Description	Species				
1	Current year seedling					
2	< 1 m (3 ft) in height					
3	> 1 m (3 ft) in height and < 10 cm (4 in) DBH					
4	>10 cm (4 in) DBH					

Site Characterization

Elevational differences along transect (e.g. terraces, gradual change):

General composition of bed material in river:

Multiple channels or single channel:

Active erosion or protection on banks:

Figure C-3.—Density and Site Characterization data form

Variable	Rankings-written description and numeric score											
	Poor ===== Excellent											
Coverage and Spatial Diversity	Site permanently converted to land use not able to support native riparian vegetation, such as housing, agriculture, or concrete channel	0	No existing riparian vegetation (e.g., covered with grasses and scrub, bare ground).	0.2	Patches of monotypic woody riparian vegetation covering up to 50% of the site, interspersed among herbaceous species or bare ground.	0.4	Patches of diverse riparian vegetation (e.g., at least two different genera of woody riparian vegetation present) covering up to 30% of the site, interspersed among grasses, invasive plants, or bare ground; and/or greater than 50% of the site covered with monotypic patches of riparian vegetation, interspersed among herbaceous species or bare ground.	0.6	Diverse woody riparian vegetation (at least three genera) covering between 30% and 75% of the site, e.g., strips or islands of riparian habitat interspersed in open space.	0.8	Diverse riparian vegetation (e.g., at least three different genera of native riparian vegetation present) covering between 75% and 100% of the site, interspersed in open space or herbaceous plant communities.	1.0
Structural Diversity	Site permanently converted to land use not able to support native riparian vegetation, such as housing, agriculture, or concrete channel	0	No existing riparian vegetation (e.g., covered with grasses and scrub, bare ground).	0.2	Vegetated areas of the site contain sparse, scattered, patchy, or remnant riparian vegetation that is immature and/or lacks structural (vertical) diversity.	0.4	Patches of riparian vegetation contain riparian trees and/or saplings(i.e., perennial dicots), but contain none or poorly developed shrub understory.	0.6	Riparian vegetation patches contain cottonwood trees and saplings, with well-developed native shrub understory, or shrub understory, but few riparian trees.	0.8	Patches of diverse riparian vegetation. They contain cottonwood trees, saplings, and seedlings (or evidence of seedling establishment), as well as developed native shrub understory and herbaceous layer.	1.0

Contiguity of Habitats	No linear contiguity or transitional upland habitat; surrounded by or isolated within an anthropogenic modified setting.	0	No linear contiguity upstream or downstream, but isolated within upland open space habitat.	0.2	Contiguous with comparable habitat on one end of the site, but surrounded with urban/suburban or other non-open space lands adjacent (lateral to) to the site on at least one side.	0.4	Contiguous with comparable habitat on one end of the site and surrounded by transitional upland habitat which is at least twice the width of the riparian zone.	0.6	Contiguous with comparable habitat on both ends of the site, but surrounded with anthropogenically modified lands adjacent (lateral to) to the site on at least one side.	0.8	Contiguous with comparable habitat on both ends of the site and surrounded by transitional upland habitat on both sides which is at least twice the width of the riparian zones.	1.0
Percent of Invasive Woody Vegetation (please note other invasive herbaceous vegetation)	Site is covered by pure stands of invasive vegetation or lacks any riparian vegetation	0	70-99% invasive vegetation.	0.2	40-69% invasive vegetation.	0.4	10-39% invasive vegetation.	0.6	4-9% invasive vegetation	0.8	Site is covered by less than 5% invasive vegetation.	1.0
Hydrology	No regular supply of water to the site. Site not associated with any water source, surface drainage, impoundment, or groundwater discharge.	0	Water supply to the site is solely from artificial irrigation. No natural supply.	0.2	Site is sustained by source of water not associated with water way. For example, the site is sustained by groundwater or urban runoff. There is no evidence of riparian processes.	0.4	Site is sustained by natural source, but no evidence of riparian processes, such as overbank flow or scour or deposition. Cut banks.	0.6	Site is within or adjacent to an impoundment on a natural waterway which is subject to fluctuations in flow or hydroperiod.	0.8	Site is within or adjacent to a waterway that provides the primary source of water to the site. The site contains evidence of riparian processes where water flows into the riparian vegetation zone.	1.0
Micro- and Macrotopographic Complexity	Flood-prone area contained in a concrete-lined channel.	0	Flood-prone area is characterized by a homogenous, flat earthen surface with little to no micro- and macro-topographic	0.2	Flood-prone area contains micro- and/or macro-topographic features such as pits, ponds, hummocks, bars, but is predominantly	0.5	Flood plain mostly heterogeneous, characterized by micro-topographic features ie pits, ponds, hummocks, bars. However, there are no macro-	0.8	Flood-prone area is characterized by micro- and macro-topographic complexity, such as meanders, bars, braiding,	1.0		

			features.		homogenous or flat surface		topographic features, such as braiding, 2° channels, backwaters.		2°channels, backwaters, terraces, pits, ponds, hummocks, etc.			
Characteristics of Flood-prone Area	All flows are contained in a concrete-lined channel, culvert, etc.	0	Channel has an earthen bottom; however, it is structurally confined (e.g., riprap or concrete sideslopes) such that the flood-prone area is wholly contained within the channel, except in extreme events.	0.2	Channel has an earthen bottom and earthen sideslopes; however, it is incised or confined such that the flood-prone area is wholly contained within the channel and there is no opportunity for overbank flow, except in extreme events.	0.3	Site is part of a flood plain, which provides an opportunity for overbank flow during moderate flow events (i.e., during a 2- to 10-year-flood event). However, the flood-prone area is confined by levees, berms, dikes, cut banks, or other obstructions or barriers such that the area available for overbank flow is less than twice the width of the channel at bankfull conditions.	0.6	Site is part of a flood plain, which provides an opportunity for overbank flow during moderate flow events. The flood-prone area is confined by levees, berms, dikes, cut banks, or other obstructions or barriers; however, the area available for overbank flow is equal to or greater than twice the width of the channel at bankfull conditions.	0.8	Site is part of an unconfined natural floodplain at least twice the width of the channel at bankfull conditions and there is evidence of overbank flow.	1.0
Biogeochemical Processes	Flood-prone area contained in a concrete-lined channel, culvert, etc., with little to no vegetation or detritus.	0	Site can support grasses, forbs, or other herbaceous vegetation, and there is woody debris, leaf litter, or detritus present in the channel.	0.2	Site supports at least 25% relative cover of grasses, forbs, herbaceous, or riparian vegetation, and there is at least 10% relative cover of woody debris, leaf litter, or detritus in the channel.	0.4	Site contains between 25% and 50% relative cover of any strata of riparian vegetation and between 10% and 40% relative cover with woody debris, leaf litter, or detritus.	0.6	Site contains between 50% and 75% relative cover of any strata of riparian vegetation and between 40% and 60% relative cover with woody debris, leaf litter, or detritus.	0.8	Site contains greater than 75% relative cover of any strata of riparian vegetation (native or non-native) and greater than 60% relative cover with woody debris, leaf litter, or detritus.	1.0

Figure C-4.—Habitat variables data form.

Appendix D

Scientific Names and Locations of Plants Detected in Vegetation Transects
2011 to 2013

CODE	SCIENTIFIC NAME	COMMON NAME	LIFEFORM	REACH										
				1A	1B	2A	2B	3	4A	ESB	MB	4B2	5	
Tree/shrub														
ALRH	<i>Alnus rhombifolia</i>	White alder	NT	X										
CEOC	<i>Cephalanthus occidentalis</i>	Button bush	NS	X	X									
FRLA	<i>Fraxinus latifolia</i>	Oregon ash	NT	X										
POFR	<i>Populus fremontii</i>	Fremont cottonwood	NT		X		X	X						
QULO	<i>Quercus lobata</i>	Valley oak	NT	X										
SAEX	<i>Salix exigua</i>	Sandbar willow	NS	X	X		X							
SAGO	<i>Salix gooddingii</i>	Gooding's willow	NT		X			X	X	X			X	X
SALAE	<i>Salix laevigata</i>	Red willow	NT			X	X							
SALAS	<i>Salix lasiolepis</i>	Arroyo willow	NT	X										
SALU	<i>Salix lucida</i>	Shining willow	NT	X										
SANI	<i>Sambucus nigra</i>	Black elderberry	NT				X							
SEPU	<i>Sesbania pungens</i>	Scarlet wisteria	IS		X									
Graminoid														
ALSA	<i>Alopecurus saccatus</i>	Pacific foxtail	NG								X		X	
ARDO2	<i>Arundo donax</i>	Giant reed	IG		X									
BRDI	<i>Bromus diandrus</i>	Ripgut brome	IG	X		X		X			X		X	
BRHO	<i>Bromus hordeaceus</i>	Soft chess brome	IG	X							X		X	
BRIN	<i>Bromus inermis</i>	Smooth brome	IG					X						
BRMA	<i>Bromus madritensis</i>	Foxtail chess	IG	X	X	X	X	X	X	X				
CYAC	<i>Cyperus acuminatus</i>	Tapertip flatsedge	NG				X			X				
CYDA	<i>Cynodon dactylon</i>	Bermuda grass	IG	X	X	X		X	X	X	X	X	X	X
CYES	<i>Cyperus esculentus</i>	Yellow nutgrass	NG		X			X	X	X	X			
CYSP	<i>Cyperus sp.</i>	Flatsedge		X										
DISP	<i>Distichlis spicata</i>	Salt grass	NG	X					X	X	X	X	X	X
ECCR	<i>Echinochloa crus-galli</i>	Barnyard grass	IG	X		X		X	X	X			X	
ELMA	<i>Eleocharis macrostachya</i>	Common spikerush	NG							X				
HOMA	<i>Hordeum marinum ssp gussoneanum</i>	Mediterranean barley	IG							X	X	X		
HOMU	<i>Hordeum murinum</i>	Foxtail barley	IG							X	X			
JUAC	<i>Juncus acuminatus</i>	Tapertip rush	NG	X										
JUBA	<i>Juncus balticus</i>	Baltic rush	NG	X			X			X	X	X	X	X
LEOR	<i>Leersia oryzoides</i>	Rice cutgrass	NG	X										
LEUN	<i>Leptochloa uninervia</i>	Mexican sprangletop	NG	X		X	X	X	X	X				
LETR	<i>Leymus triticoides</i>	Creeping wildrye	NG							X				
MUAS	<i>Muhlenbergia asperifolia</i>	Scratchgrass	NG					X						
ORSA	<i>Oryza sativa</i>	Rice	IG			X								
PADI	<i>Paspalum dilatatum</i>	Dallis grass	IG	X		X				X	X			
PANO	<i>Paspalum notatum</i>	Bahia grass	IG							X				
PHAR	<i>Phalaris arundinacea</i>	Canary reedgrass	NG	X										
POMO	<i>Polypogon monspeliensis</i>	Rabbitsfoot grass	IG							X				X
VUMY	<i>Vulpia myuros</i>	Rat-tail fescue	IG	X	X	X	X	X		X				
Forb														
AMRO	<i>Ammania robusta</i>	Grand redstem	NF							X	X	X		
ANCO	<i>Anthemis cotula</i>	Dog fennel	IF							X				
ARDO	<i>Artemisia douglasiana</i>	California mugwort	NF	X	X		X	X					X	X
ARLU	<i>Artemisia ludoviciana</i>	White sagebrush	NF	X										
ARVU	<i>Artemisia vulgare</i>	Common mugwort	IF			X			X					
BRNI	<i>Brassica nigra</i>	Black mustard	IF	X		X	X		X	X	X	X	X	X
CESO	<i>Centaurea solstitialis</i>	Yellow starthistle	IF							X				

CODE	SCIENTIFIC NAME	COMMON NAME	LIFEFORM	REACH									
				1A	1B	2A	2B	3	4A	ESB	MB	4B2	5
CEPA	<i>Centromadia parryii ssp rudis</i>	Pappose tarweed	NF							X	X	X	
CHCA	<i>Chenopodium californicum</i>	California goosefoot	NF					X			X	X	X
CHLI	<i>Chrysothamnus linifolius</i>	Spearleaf rabbitbrush	NF	X	X							X	X
CIVU	<i>Cirsium vulgare</i>	Bull thistle	IF							X	X	X	
COMA	<i>Conium maculatum</i>	Poison hemlock	IF									X	
COCA	<i>Conyza canadensis</i>	Horseweed	NF				X		X				X
CUSP	<i>Cuscuta sp.</i>	Dodder	IF				X						
DAWR	<i>Datura wrightii</i>	Jimson weed	NF										X
ERSE	<i>Eremocarpus setigerus</i>	Doveweed	NF			X							
ERWR	<i>Eriogonum wrightii</i>	Wright's buckwheat	NF			X							
ERCI	<i>Erodium cicutarium</i>	Redstem storks bill	IF				X		X				
ERRA	<i>Eryngium racemosum</i>	Delta button celery	NF							X			
ESCA	<i>Eschscholzia californica</i>	California poppy	NF		X	X							
GATR	<i>Gallium trifudum</i>	Threepetal bedstraw	NF					X					
GRCA	<i>Grindelia camporum</i>	Gum plant	NF							X	X	X	X
HEAN	<i>Helianthus annuus</i>	Sunflower	NF			X		X			X	X	
HECU	<i>Heliotropium curassavicum</i>	Salt heliotrope	NF								X		X
KOSC	<i>Kochia scoparia</i>	Kochia	IF			X	X	X		X			
LASE	<i>Lactuca serriola</i>	Prickly lettuce	IF			X		X		X	X	X	X
LELA	<i>Lepidium latifolium</i>	Perennial peppergrass	IF									X	X
LEPU	<i>Ludwigia peploides</i>	Water primrose	NF								X		
LOCO	<i>Lotus corniculatus</i>	Birdsfoot trefoil	IF							X		X	
LOUN	<i>Lotus unifoliolatus</i>	American bird's-foot trefoil	NF			X	X						
MALE	<i>Malvella leprosa</i>	Alkali mallow	IF							X		X	X
MEAL	<i>Melilotus alba</i>	White sweetclover	IF						X	X			X
MEAR	<i>Mentha arvensis</i>	Field mint	NF	X			X						
MEPU	<i>Mentha pulegium</i>	Pennyroyal	NF	X									
MYAQ	<i>Myriophyllum aquaticum</i>	Parrotfeather	IF	X									
PHNO	<i>Phyla nodiflora</i>	Turkey tangle fogfruit	NF							X	X	X	
POAR	<i>Polygonum arenastrum</i>	Common knotweed	IF							X	X	X	
POLA	<i>Polygonum lapathifolium</i>	Pale smartweed	NF					X					
PSCA	<i>Pseudognaphalium californicum</i>	California cudweed	NF	X	X	X	X			X			X
ROPA	<i>Rorippa palustris</i>	Yellow cress	NF				X	X	X			X	X
RUCR	<i>Rumex crispus</i>	Curly dock	IF					X	X	X	X	X	
RUDI	<i>Rubus discolor</i>	Himalayan blackberry	IF	X									
SATR	<i>Salsola tragus</i>	Russian thistle	IF			X							
SASP	<i>Salsola sp.</i>	Saltwort	IF					X	X	X			
SIMA	<i>Silybum marianum</i>	Milk thistle	IF							X			
SOAM	<i>Solanum americanum</i>	American black nightshade	NF			X	X					X	X
SOAS	<i>Sonchus asper</i>	Prickly sow thistle	IF				X			X			
TRSP	<i>Trifolium sp.</i>	Clover					X	X		X			
URDI	<i>Urtica dioica</i>	Stinging nettle	IF				X						
VEAN	<i>Veronica anagallis-aquatica</i>	Water speedwell	IF		X								
XAST	<i>Xanthium strumarium</i>	Cocklebur	NF	X		X	X	X	X	X	X	X	X

Appendix E

Total percent cover of individual plant species detected
in the understory layer of vegetation transects
from 2011 to 2013.

Table E-1a. Average Total Percent Understory Cover – Upstream Reaches

Species	River Reach														
	1A			1B			2A			2B			3		
	2011	2012	2013	2011	2012	2013	2011	2012	2013	2011	2012	2013	2011	2012	2013
Button bush	0	0	0	0.5	0	0	0	0	0	0	0	0	0	0	0
Sandbar willow	0.5	0	0.5	4.0	1.5	1.0	0	0	0	0	0	0.5	0	0	0
Goodding's willow	0	0	0	0	0	1.5	0	0	0	0	0	0	0.3	0	0
Red willow	0	0	0	0	0	0	0	0	0	0.5	1.0	2.5	0	0	0
Shining willow	0	0.5	0	0	0	0	0	0	0	0	0	0	0	0	0
Oregon ash	0	0	0.5	0	0	0	0	0	0	0	0	0	0	0	0
Native trees/shrubs	0.5	0.5	1.0	4.5	1.5	2.5	0	0	0	0.5	1.0	3.0	0.3	0	0
Scarlet wisteria	0	0	0	0	0.5	0	0	0	0	0	0	0	0	0	0
Introduced trees/shrubs	0	0	0	0	0.5	0	0	0	0	0	0	0	0	0	0
Yellow nutgrass	0	0	0	5.0	0	0	0	0	0	0	0	0	3.5	0	0
Flatsedge	1.0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Salt grass	0.5	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Baltic rush	1.5	0	1.0	0	0	0	0	0	0	0	0	1.5	0	0	0
Mexican sprangletop	0.5	0	0	0	0	0	1.5	0.5	0	2.2	0	0	1.6	0	0
Canary reedgrass	0	0.5	0	0	0	0	0	0	0	0	0	0	0	0	0
Tapertip rush	0	0	1.0	0	0	0	0	0	0	0	0	1.2	0	0	0
Rice cutgrass	0	0	3.0	0	0	0	0	0	0	0	0	0	0	0	0
Scratchgrass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.7
Unidentified grasses*	0.5	2.5	0	0	0	0	0	0	0	0	0	0	0	0	0
Native graminoids	4.0	3.0	5.0	5.0	0	0	1.5	0.5	0	2.2	0	2.7	5.1	0	0.7
Ripgut brome	0.5	13.0	22.5	0	0	0	3.0	6.0	6.0	0	0	0	0	16.0	35.5
Bermuda grass	1.0	1.5	0.5	2.0	10.5	4.0	0.0	0.5	1.0	0	0	0	4.1	1.6	0.7
Barnyard grass	0.5	0	0	0	0	0	0.5	0	0	0	0	0	0.6	0	0
Rice	0	0	0	0	0	0	1.0	0	0	0	0	0	0	0	0
Soft chess brome	0	3.0	1.5	0	0	0	0	0	0	0	0	0	0	0	0
Foxtail chess	0	3.0	0	0	5.0	12.0	0	4.0	12.0	0	0	3.4	0	0	0.3
Dallis grass	0	0	1.0	0	0.0	0	1.0	0	0	0	0	0	0	0	0
Rat-tail fescue	0	10.0	8.5	0	0.5	0	1.5	2.0	2.5	0	0	0.7	0	0	4.2
Smooth brome	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.3
Giant reed	0	0	0	0	0	0.5	0	0	0	0	0	0	0	0	0
Introduced graminoids	2.0	30.5	34.0	2.0	16.0	16.5	7.0	12.5	21.5	0.0	0.0	4.1	4.7	17.6	41.0
California mugwort	8.0	4.5	6.5	0	0	0.5	0	0	0	1.4	2.9	2.1	10.5	13.6	14.8
California goosefoot	0	0	0	0	0	0	0	0	0	0	0	0	3.3	0	0
Doveweed	0	0	0	0	0	0	0.5	0	0	0	0	0	0	0	0
Wright's buckwheat	0	0	0	0	0	0	1.5	0.5	0	0	0	0	0	0	0
Threepetal bedstraw	0	0	0	0	0	0	0	0	0	0	0	0	2.5	0	0
Sunflower	0	0	0	0	0	0	1.0	0	0	0	0	0	0	0	0.9
American bird's-foot trefoil	0	0	0	0	0	0	3.5	0	0.5	9.3	0	4.3	0	0	0
Field mint	0.5	1.0	0	0	0	0	0	0	0	0	0	0.7	0	0	0
Pale smartweed	0	0	0	0	0	0	0	0	0	0	0	0	12.4	0	0
Yellow cress	0	0	0	0	0	0	0	0	0	0	0.7	0	2.2	0	1.3
American black nightshade	0	0	0	0	0	0	0.5	0	0	4.3	0	0	2.8	0	0
Cocklebur	1.0	0	0	0	0	0	0.5	0	0	0	0	1.2	20.6	0	0
Spearleaf rabbitbrush	0	0.5	1.5	0	1.5	6.5	0	0	0	0	0	0	0	0	0
Pennyroyal	0	0.5	0	0	0	0	0	0	0	0	0	0	0	0	0
California poppy	0	0	0	0	0.5	0.5	0	1.0	2.0	0	0	0	0	0	0
California cudweed	0	0	1.0	0	0.5	0	0	0.5	0	0	0.5	0.7	0	0	0
Horseweed	0	0	0	0	0	0	0	0	0	0	1.2	0	0	0	0
White sagebrush	0	0	0.5	0	0	0	0	0	0	0	0	0	0	0	0
Unidentified forbs*	0.5	0	0	0	0	0.5	0	0	0	1.4	0	0	0.7	0	0
Native forbs	10.0	6.5	9.5	0	2.5	8.0	7.5	2.0	2.5	16.4	5.3	9.0	55.0	13.6	17.0
Black mustard	1.0	0	0	0	0	0	2.0	4.5	1.5	4.1	12.4	7.2	0	0	0
Prickly lettuce	0	0	0	0	0	0	0.5	0	0	0	0	0	0.8	0	0
Curly dock	0	0	0	0	0	0	0	0	0	0	0	0	0	0.3	0.3
Himalayan blackberry	5.5	8.0	5.0	0	0	0	0	0	0	0	0	0	0	0	0
Russian thistle	0	0	0	0	0	0	0.5	0	0	0	0	0	0	0	0
Saltwort	0	0	0	0	0	0	0	0	0	0	0	0	7.5	0	0

Table E-1a. Average Total Percent Understory Cover – Upstream Reaches															
Species	River Reach														
	1A			1B			2A			2B			3		
	2011	2012	2013	2011	2012	2013	2011	2012	2013	2011	2012	2013	2011	2012	2013
Clover	0	0	0	0	0	0	0	0	0	2.1	4.3	0	0	0	0.9
Stinging nettle	0	0	0	0	0	0	0	0	0	0.5	0.5	0.5	0	0	0
Redstem storks bill	0	0	0	0	0	0	0	0	0	0	0.7	0	0	0	0
Koschia	0	0	0	0	0	0	0	0.5	0	0	0.7	0	0	0	0.9
Prickly sowthistle	0	0	0	0	0	0	0	0	0	0	0.7	0	0	0	0
Water speedwell	0	0	0	0.5	0	0	0	0	0	0	0	0	0	0	0
Dodder	0	0	0	0	0	0	0	0	0	0	0	2.5	0	0	0
Common mugwort	0	0	0	0	0	0	0	0	1.0	0	0	0	0	0	0
Parrotfeather	0	0	1.0	0.5	0	0	0	0	0	0	0	0	0	0	0
Introduced forbs	6.5	8.0	6.0	1.0	0	0	3.0	5.0	2.5	6.7	19.3	10.2	8.3	0.3	2.1
Total Plant Cover	23.0	48.5	55.5	12.5	20.5	27.0	19.0	20.0	26.5	25.8	25.6	29.0	73.4	31.5	60.8
Litter	33.5	39.0	39.0	20.0	40.5	42.5	16.5	17.5	15.5	22.9	32.8	33.3	16.0	56.9	34.5
Bare	37.0	6.0	2.5	55.0	32.5	30.5	45.0	43.0	58.0	51.3	41.6	37.8	10.6	11.6	4.8
Rock	6.0	5.0	2.5	13.0	6.5	0	19.5	19.5	0	0	0	0	0	0	0
Water	0.5	1.5	0.5	0	0	0	0	0	0	0	0	0	0	0	0
Total Cover	100.0	100.0	100.0	100.5	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.1	100.0	100.0	100.1

*Unidentified species may be either native or introduced

Table E-1b. Average Total Percent Understory Cover – Downstream Reaches													
Species	River Reach												
	4A*		ESB			MB			4B2			5	
	2011	2012	2011	2012	2013	2011	2012	2013	2011	2012	2013	2012	2013
Goodding's willow	1.5	0.7	0	0	0	0	0	0	0	0	0	0.5	0
Native trees/shrubs	1.5	0.7	0	0	0	0	0	0	0	0	0	0.5	0
Pacific foxtail	0	0	0	0	0	5.5	0	0	1.0	0	0	0	0
Tapertip flatsedge	0	0	1.0	2.3	0.3	0	0	0	0	0	0	0	0
Yellow nutgrass	1.4	0	1.0	0	0	0.5	0	0	0	0	0	0	0
Salt grass	0	4.9	0	10.7	9.5	0.0	1.5	5.0	14.4	19.1	22.6	0.0	0.5
Common spikerush	0	0	11.0	0	0	0	0	0	0	0	0	0	0
Baltic rush	0	0	0.7	0	0	6.5	6.5	3.5	5.2	5.1	5.7	0	0.5
Mexican sprangletop	2.2	0	0.2	0	0	0	0	0	0	0	0	0	0
Creeping wildrye	0	0	0	1.0	1.5	0	0	0	0	0	0	0	0
Unidentified grasses*	0	0	2.0	0	0	0	0	0	0	0	0	0	0
Native graminoids	3.6	4.9	15.9	14.0	11.3	12.5	8.0	8.5	20.6	24.2	28.3	0.0	1.0
Bermuda grass	1.2	0	2.7	3.0	0	3.0	0	0	2.0	4.0	3.5	1.0	0
Barnyard grass	1.7	0	0.3	0	0	0	0	0	26.1	0	0	0	0
Mediterranean barley	0	0	0	0	0.3	5.0	5.5	14.5	1.0	2.5	1.0	0	0
Bahia grass	0	0	5.3	0	0	0	0	0	0	0	0	0	0
Soft chess brome	0	0	0	0	0	0	2.0	0	0	5.5	11.2	0	0
Foxtail chess	0	3.2	0	0.3	0.5	0	0	0	0	0	0	0	0
Rabbitsfoot grass	0	0	0	1.4	0.8	0	0	0	0	0	0	5.5	2.0
Foxtail barley	0	0	0	0.3	0	0	2.0	7.0	0	0	0	0	0
Dallis grass	0	0	4.5	1.7	3.5	0	0	1.5	0	0	0	0	0
Rat-tail fescue	0	0	0	0.3	0	0	0	0	0	0	0.5	0	0
Rippgut brome	0	0	0	0.3	0	0	0	1.5	0	0	3.8	0	0
Introduced graminoids	2.9	3.2	12.8	7.3	5.1	8.0	9.5	24.5	29.1	12.0	20.0	6.5	2.0
California mugwort	0	0	0	0	0	0	0	0	0	1.6	0.6	0.5	4.5
Pappose tarweed	0	0	0	0	2.5	33.0	0	1.0	1.3	0	0	0	0
California goosefoot	0	0	0	0	0	1.5	0	0	3.5	1.3	4.0	0.5	0
Delta button celery	0	0	0.5	0	0	0	0	0	0	0	0	0	0
Sunflower	0	0	0	0	0	3.0	0	0	7.1	0	0	0	0
Yellow cress	0	0.5	0	0	0	0	0	0	0	0.6	0	2.5	0
American black nightshade	0	0	0	0	0	0	0	0	1.9	0	0	1.5	0
Cocklebur	4.2	0	7.3	2.7	0.3	12.0	0	0	7.1	0	0	1.0	0
Cottonbatting cudweed	0	0	0	0.5	0	0	0	0	0	0	0	2.0	0
Horseweed	0	0.5	0	0	0	0	0	0	0	0	0	1.0	0.5
Gumweed	0	0	0	0.3	1.7	0	4.5	6.5	0	1.1	6.2	0	2.0
Turkey tangle fogfruit	0	0	0	5.2	4.7	0	4.0	8.5	0	0	0.5	0	0

Table E-1b. Average Total Percent Understory Cover – Downstream Reaches

Species	River Reach												
	4A*		ESB			MB			4B2			5	
	2011	2012	2011	2012	2013	2011	2012	2013	2011	2012	2013	2012	2013
Jimson weed	0	0	0	0	0	0	0	0	0	0	0	1.0	1.0
Grand redstem	0	0	3.0	0	0	0	0.5	0	0	0	0	0	0
Spearleaf rabbitbrush	0	0	0	0	0	0	0	0	0	0	0.6	0	1.5
Water primrose	0	0	0	0	0.5	0	0	0	0	0	0	0	0
Salt heliotrope	0	0	0	0	0	0	0	0.5	0	0	0	0	0.5
Unidentified forbs*	2.0	0	3.0	2.5	0.5	1.5	0	0	0	0	0	9.0	0
Native forbs	6.2	1.0	13.8	11.2	10.2	51.0	9.0	16.5	20.9	4.6	11.9	19.0	10.0
Common mugwort	2.9	0	0	0	0	0	0	0	0	0	0	0	0
Black mustard	0	4.6	0	1.3	2.7	9.0	16.5	2.0	5.8	10.3	5.8	1.0	3.5
Prickly lettuce	0	0	0	0.3	1.0	1.0	0	0	1.0	0	0	1.0	0
Alkali mallow	0	0	1.0	1.3	0.5	0	0	0	0.6	0	0	1.0	0
Common knotweed	0	0	0	0.5	0	9.5	0	0	2.1	1.9	0	0	0
Curly dock	0.7	0	0.5	3.0	0	3.0	0	0	1.3	0.6	0	0	0
Saltwort	0.5	0	0	0	0	0	0	0	0	0	0	0	0
Clover	0	0	3.3	0	0	0	0	0	0	0	0	0	0
Redstem storks bill	0	0.5	0	0	0	0	0	0	0	0	0	0	0
Prickly sowthistle	0	0	0	0.7	0	0	0	0	0	0	0	0	0
White sweetclover	0	5.7	0	0.5	0	0	0	0	0	0	0	9.0	6.0
Bull thistle	0	0	0	0	1.8	0	0.5	0	0	1.3	1.3	0	0
Poison hemlock	0	0	0	0	0	0	0	0	0	9.3	1.9	0	0
Perennial pepperweed	0	0	0	0	0	0	0	0	0	1.9	2.5	7.5	22.5
Birdsfoot trefoil	0	0	0	0.5	1.5	0	0	0	0	0.6	0	0	0
Dog fennel	0	0	0	2.7	0	0	0	0	0	0	0	0	0
Milk thistle	0	0	0	0.7	0	0	0	0	0	0	0	0	0
Kochia	0	0	0	0	0.5	0	0	0	0	0	0	0	0
Yellow starthistle	0	0	0	0	2.5	0	0	0	0	0	0	0	0
Introduced forbs	4.1	10.8	4.8	11.5	10.5	22.5	17.0	2.0	10.8	25.9	11.5	19.5	32.0
Total Plant Cover	18.3	20.6	47.3	44.0	37.1	94.0	43.5	51.5	81.4	66.7	71.7	45.5	45.0
Litter	7.5	26.7	22.7	39.5	49.3	4.5	55.0	44.0	11.8	32.8	28.4	42.0	49.0
Bare	74.2	52.7	30.0	16.8	13.8	1.5	1.5	4.5	6.8	0.5	0.0	12.5	6.0
Total Cover	100.0	100.0	100.0	100.3	100.2	100.0	100.0	100.0	100.0	100.0	100.1	100.0	100.0

*4A was not sampled in 2013

Appendix F

Photo Stations
August 2011 and June 2013

Reach 1A, Transect 1

1a – Toward transect



August 2011



June 2012



June 2013

1a – Away from transect



August 2011



June 2012



June 2013

1b – Toward transect



August 2011



June 2012



June 2013

1b – Away from transect



August 2011



June 2012



June 2013

Reach 1A, Transect 2

2a – Toward transect



August 2011



June 2012



June 2013

2a – Away from transect



August 2011



June 2012



June 2013

2b – Toward transect



August 2011



June 2012



June 2013

2b – Away from transect



August 2011



June 2012



June 2013

Reach 1B, Transect 1

1a – Toward transect



August 2011



June 2012



June 2013

1a – Away from transect



August 2011



June 2012



June 2013

1b – Toward transect



August 2011



June 2012



June 2013

1b – Away from transect (tape continues beyond transect)



August 2011



June 2012



June 2013

Reach 1B, Transect 2

2a – Toward transect



August 2011



June 2012



June 2013

2a – Away from transect



August 2011



June 2012



June 2013

2b – Toward transect



August 2011



June 2012



June 2013

2b – Away from transect



August 2011



June 2012



June 2013

Reach 2A, Transect 1

1a – Toward transect



August 2011



June 2012



June 2013

1a – Away from transect



August 2011




June 2012



June 2013

1b – Toward transect



August 2011



June 2012



June 2013

1b – Away from transect



August 2011



June 2012



June 2013

Reach 2A, Transect 2

2a – Toward transect



August 2011



June 2012



June 2013

2a – Away from transect



August 2011



June 2012



June 2013

2b – Toward transect



August 2011



June 2012



June 2013

2b – Away from transect



August 2011




June 2012



June 2013

Reach 2B, Transect 1


1a – Toward transect



August 2011



June 2012



June 2013

1a – Away from transect



August 2011



June 2012



June 2013

1b – Toward transect



August 2011



June 2012



June 2013

1b – Away from transect



August 2011




June 2012



June 2013

Reach 2B, Transect 2

2a – Toward transect



August 2011



June 2012



June 2013

2a – Away from transect



August 2011



June 2012



June 2013

2b – Toward transect (taken from different angle in 2011)



August 2011



June 2012



June 2013

2b – Away from transect



August 2011



June 2012



June 2013

Reach 3, Transect 1

1a – Toward transect



August 2011



June 2012



June 2013

1a – Away from transect



August 2011



June 2012



June 2013

1b – Toward transect



August 2011



June 2012



June 2013

1b – Away from transect



August 2011



June 2012



June 2013

Reach 3, Transect 2

2a – Toward transect



August 2011



June 2012



June 2013

2a – Away from transect



August 2011



June 2012



June 2013

2b – Toward transect



August 2011



June 2012



June 2013

2b – Away from transect



August 2011




June 2012



June 2013

Reach 4A, Transect 1

1a – Toward transect



August 2011



June 2012

1a – Away from transect



August 2011



June 2012

1b – Toward transect



August 2011



June 2012

1b – Away from transect



August 2011



June 2012

Reach 4A, Transect 2

2a – Toward transect



August 2011



June 2012

2a – Away from transect



August 2011



June 2012

2b – Toward transect



August 2011



June 2012

2b – Away from transect



August 2011



June 2012

East Side Bypass, Transect 1


1a – Toward transect



August 2011



June 2012



June 2013

1a – Away from transect



August 2011




June 2012



June 2013


1b – Toward transect



August 2011




June 2012



June 2013


1b – Away from transect



August 2011



June 2012



June 2013

East Side Bypass, Transect 2

2a – Toward transect



August 2011



June 2012



June 2013

2a – Away from transect



August 2011



June 2012



June 2013


2b – Toward transect



August 2011



June 2012



June 2013


2b – Away from transect



August 2011



June 2012



June 2013

East Side Bypass (Merced NWR), Transect 3


3a – Toward transect



August 2011



June 2012



June 2013

3a – Away from transect



August 2011



June 2012




June 2013

3b – Toward transect



August 2011



June 2012



June 2013

3b – Away from transect



August 2011



June 2012



June 2013

East Side Bypass (Merced NWR), Transect 4

4a – Toward transect



August 2011



June 2012



June 2013

4a – Away from transect



August 2011



June 2012



June 2013

4b – Toward transect



August 2011



June 2012




June 2013

4b – Away from transect



August 2011



June 2012




June 2013

Mariposa Bypass, Transect 1

1a – Toward transect



August 2011



June 2012



June 2013

1a – Away from transect



August 2011



June 2012



June 2013

1b – Toward transect



August 2011



June 2012



June 2013

1b – Away from transect



August 2011



June 2012



June 2013

Mariposa Bypass, Transect 2

2a – Toward transect



August 2011



June 2012



June 2013

2a – Away from transect



August 2011



June 2012



June 2013

2b – Toward transect



August 2011



June 2012




June 2013

2b – Away from transect



August 2011



June 2012



June 2013

Reach 4B2 (San Luis NWR), Transect 1

1a – Toward transect



August 2011



June 2012



June 2013

1a – Away from transect



August 2011



June 2012



June 2013

1b – Toward transect



August 2011



June 2012



June 2013

1b – Away from transect



August 2011



June 2012



June 2013

Reach 4B2 (San Luis NWR), Transect 2

2a – Toward transect



August 2011



June 2012



June 2013

2a – Away from transect



August 2011



June 2012



June 2013

2b – Toward transect



August 2011



June 2012



June 2013

2b – Away from transect



August 2011



June 2012



June 2013

Reach 5, Transect 1

1a – Toward transect



June 2012



June 2013

1a – Away from transect



June 2012



June 2013

1b – Toward transect



June 2012



June 2013

1b – Away from transect



June 2012



June 2013

Reach 5, Transect 2

2a – Toward transect



June 2012



June 2013

2a – Away from transect



June 2012



June 2013

2b – Toward transect



June 2012



June 2013

2b – Away from transect



June 2012



June 2013

PEER REVIEW DOCUMENTATION

PROJECT AND DOCUMENT INFORMATION

Project Name San Joaquin River Restoration Project
WOID A207F
Document Vegetation Response to Interim Flows in the San Joaquin River
Document Date April 2014
Team Leader Gregory Reed
Document Author(s)/Preparer(s) Rebecca Siegle, Gregory Reed, S. Mark Nelson, Erin Rice
Peer Reviewer Gregory Reed
Peer Reviewer S. Mark Nelson

REVIEW REQUIREMENT

Part A: Document Does Not Require Peer Review



Explain _____

Part B: Document Requires Peer Review: SCOPE OF PEER REVIEW

Peer Review restricted to the following items/Section(s): _____ Reviewer: _____

REVIEW CERTIFICATION

Peer Reviewer - I have reviewed the assigned items/Section(s) noted for the above document and believe them to be in accordance with the project requirements, standards of the profession, and Reclamation policy.

Reviewer: Gregory Reed Review Date: 1-28-14 Signature: 
Reviewer: S. Mark Nelson Review Date: 1-28-14 Signature: 

I have discussed the above document and review requirements with the Peer Reviewer and believe that this review is completed, and that the document will meet the requirements of the project.

Team Leader: Gregory Reed Date: 4-7-14 Signature: 